



# PROTECTING POST-HARVEST GRAINS FROM PROFIT-ROBBING INSECTS





# TABLE OF CONTENTS

## **Why Protect Stored Grains From Insects?**

The Economic Advantage .....	2
The Impact of Infestations .....	3

<b>Target Insect Species</b> .....	4-5
------------------------------------	-----

## **Post-Harvest Grain Storage**

Using S.L.A.M. ....	6
Sites to Treat .....	6
Insect Control and FSMA .....	7
Sidebar Feature: Life Cycle Of Grain Weevil .....	7

## **Protecting Stored Grains**

Sanitation .....	8
Treating Empty Bins and Cleaning Equipment Prior to Harvest .....	8
Treating Grains During Loading .....	8
Treating Existing Infestations .....	9
Sidebar Feature: Tips & Tricks for Treating Grain .....	9

## **Our Products**

Insect Growth Regulators .....	10
Adulticides and Synergists .....	11
Combination Products .....	12-13
Premise Control .....	13

<b>Product Application and Dilution Rates</b> .....	14-17
---	-------

<b>Specimen Labels</b> .....	18-50
------------------------------	-------

# MEET CENTRAL LIFE SCIENCES

**CENTRAL LIFE SCIENCES IS DEDICATED TO IMPROVING THE HEALTH, WELL-BEING AND THE PROTECTION** of companion animals, economic animals, plants, stored grains and the environment. To enhance the quality of life through products that are researched and developed in a scientific environment, we create trusted consumer and professionally applied goods to control economic, nuisance and disease-carrying insects throughout their life cycles.

Wheat | Corn | Oats | Rye



Rice | Sorghum | Barley | Sunflower



## LET'S PROTECT GRAINS TOGETHER

**PROTECTING STORED GRAINS FROM THE RAVAGING DAMAGE AND PROFIT LOSS CAUSED BY INSECTS HAS BEEN AN ONGOING CONCERN OF AGRICULTURAL PROFESSIONALS FOR CENTURIES.**

Procuring the right lineup of insect control procedures and products is an essential element of protection. That's why Central Life Sciences continues to develop a unique grouping of innovative products that provide protection from infestations throughout the insect life cycle that we call, "*Bug Free Grains Stored Product Solutions.*" From rescue treatments and empty-bin applications to on-grain treatments, Central Life Sciences has you covered.

Bug Free Grains Stored Product Solutions are a quality lineup of grain protectants that are formulated to control labeled targeted stored product insects. Insect growth regulators, synergists, insecticides and combination products are designed for use during the sanitation and grain storage processes, available in liquid and dry formats to allow for an array of application methods.



# WHY PROTECT STORED GRAINS?


## THE ECONOMIC ADVANTAGE



**IT IS ESTIMATED THAT THE U.S. AGRICULTURE INDUSTRY LOSES UPWARDS OF \$2.5 BILLION PER YEAR AS A RESULT OF STORED GRAINS LOST TO INSECT INFESTATIONS.** That economic loss can be felt throughout the grain storage process as insects interfere with grain quality and the ability to move that grain into the market. As a result, ag professionals sacrifice revenues and their return on investment while the country's population loses food.

Up to 10% of the U.S. grain crop gets destroyed every year because of post-harvest insect infestations. The longer commodities are held in storage, the greater the risk of stored product insect infestations. Since the commodity market determines the amount of time grain will be stored, operators need to have a system in place to ensure grain protection.

To protect the commodities as well as profits, it is essential for operators to treat grains and follow best practices for **sanitation, loading, aeration and monitoring (S.L.A.M.)**. The S.L.A.M. post-harvest integrated pest management (IPM) strategy is a systems approach to maximize grain quality. To learn more about how S.L.A.M. works, turn to the Post-Harvest Grain Storage section of this guide.



**S**ANITATION

**L**OADING

**A**ERATION

**M**ONITORING

# THE IMPACT OF INFESTATIONS



Unprotected Wheat Infested with Lesser Grain Borer



Unprotected Oats Infested with Sawtoothed Grain Beetle



*\*in a lab setting*

## LIFE CYCLE

## SIGNS OF INFESTATION



### Lesser Grain Borer

Measuring about 1/8" long, this brown/black borer has a cylindrical body and a head that is not visible from above. A voracious feeder and strong flier, it bores into grain kernels and is capable of rapidly spreading infestations. Larvae are white, c-shaped and feature three pairs of small legs.

Varies based on temperature, minimum 25 days

Live adult insects; fragmented kernels; powdery residue; appearance of "shot hole" in grain; a sweet musty odor



### Rice Weevil

This small snout beetle feeds on whole and broken kernels and can infest grain in fields. With dull red-brown or black coloring, it features markings, functional wings and a densely-pitted thorax. It measures about 3/32" - 1/8" long.

Minimum 28 days

Live adult weevils; damaged grains with "shot hole" appearance



### Indian Meal Moth

Featuring a distinctive rusty brown or bronze color on its wing tips, this small brownish moth can be found in many types of pantry foods in addition to processing and storage facilities. It measures about 3/8" long with a wingspan ranging from 1/2" to 3/4". Larvae are off-white with brown heads and measure about 2/3" long.

Minimum 42 days; adult moths do not feed and typically die within a week after laying eggs

Dense webbing produced by larvae on food fragments; flying adults present in the vicinity of grain bins; larvae climbing up walls



### Sawtoothed Grain Beetle

A secondary pest of oats, wheat, barley and animal feed, this flightless beetle has a brown, flattened body measuring 1/10" - 1/8". It features distinctive sawtooth projections on the sides of its thorax and cockroach-like running legs.

Minimum 28 days

Overheated grains; visible population build-up

## LIFE CYCLE

## SIGNS OF INFESTATION



### Confused Flour Beetle

This flightless, reddish-brown beetle feeds primarily on grain dust, broken kernels and the milled grains in flour and cereal. It measures approximately  $\frac{1}{8}$ " long and has an antenna that gradually expands towards its end.

Minimum 42 days;  
can live up to  
18 months

Live adult insects;  
foul odor and taste in  
infested product; shed  
skins and fecal matter



### Granary Weevil

This flightless weevil feeds on whole and broken kernels and has a cylindrical, shiny body that measures between  $\frac{1}{3}$ " and  $\frac{1}{8}$ ". It is cold tolerant and features a long head with a distinct snout, ridged wing-covers and a thorax marked with oval pits.

Minimum 28 days

Live adult weevils;  
damaged grains with  
"shot hole" appearance



### Red Flour Beetle

Generally classified as a scavenger, this reddish-brown beetle feeds on broken or damaged grains during its immature stages. A common pest in mills and warehouses, it measures  $\frac{1}{8}$ " long.

Egg to adult within  
30 days; can live up  
to 18 months

Live adult insects;  
pungent odor,  
nauseous taste  
and potentially pink  
coloration in infested  
foods



Insects infesting grains causing  
damage and profit loss.

## USING S-L-A-M

An IPM strategy using S.L.A.M. depends on the proper selection of crop varieties, production and harvest practices, grain handling equipment, drying systems and storage management. **THE OBJECTIVE IS TO PROTECT THE QUALITY OF GRAINS FROM WEATHER, RODENTS, INSECTS AND MOLDS THROUGH FOUR SIMPLE STEPS.**

**SANITATION** refers to keeping storage facilities clean and repaired. This can include fixing holes and cracks, cleaning out all grains from previous years and treating empty bins. This is a crucial step when it comes to starting your post-harvest storage off right.

**LOADING** must be done strategically. To improve air circulation, grain should be evenly spread as it enters the storage facility. If desired, insect control products can be distributed through the grain as it is loaded into the storage site.

**AERATION** lowers the grain's temperature and reduces moisture, which in turn slows down insect reproduction rates, mold growth and more. Aeration is necessary, as warm, moist grain is an ideal habitat for insects.

**MONITORING** your sanitation, loading, aeration and treatment techniques is essential to tracking the efficacy of your IPM plan. Regularly probe and screen grains, and set up insect traps to identify pests. Techniques should be adjusted based on monitoring.

**Bug Free Grains Stored Product Solutions can go anywhere grains are to be stored.**



Silos

Grain Elevators

Warehouses

Storage Bins

Ground Piles

# INSECT CONTROL AND FSMA

The Food Safety Modernization Act (FSMA) has established Current Good Manufacturing Practices (CGMPs) for both human food and animal food production. According to official regulations, all covered facilities must implement a food safety system that includes the following:

## HAZARD ANALYSIS:

Identification of foreseeable chemical, biological and physical hazards, including injurious insects.

## PREVENTATIVE CONTROLS:

Measures that ensure the prevention or minimization of identified hazards.

## OVERSIGHT AND MANAGEMENT OF PREVENTATIVE CONTROLS:

Monitoring the efficacy of preventative controls, verifying control methods and creating a recall plan.



## UNDERSTANDING A PROBLEM PEST: GRAIN WEEVILS

Because of the way weevils transform from an egg to an adult isolated inside the grain kernel, they can be a difficult insect to control. As weevils proliferate across the U.S., identifying, preventing and managing weevil infestations all contribute to maintaining the quality and profitability of stored grains.

## GRAIN WEEVIL LIFE CYCLE



## WEEVILS IN YOUR AREA? WE HAVE THE RIGHT SOLUTIONS.



**Pre-bin treatment:**  
Control insects, larvae and eggs you can't see.

Diacon® IGR PLUS tank-mixed with PBO-8® Synergist



**Short-term storage (less than 6 months):**  
Use as a rescue treatment.

Centynal™ Synergized Insecticide



**Long-term storage (6 to 12 months):**  
Use as a rescue treatment.

Gravista® Insecticide  
Gravista®-D Insecticide

If an existing weevil infestation is present, consider treatment with an EPA-registered fumigant.



# PROTECTING STORED GRAINS

## FORMULAS FOR SUCCESSFULLY PROTECTING STORED GRAINS FROM DAMAGING INSECTS

### SANITATION

It is important to take a sample as grain comes in and hold it to see if insects emerge due to field infestations. The success of an IPM program that includes S.L.A.M. is largely dependent on steps taken before grain is even loaded into storage.

### TREATING EMPTY BINS AND CLEANING EQUIPMENT PRIOR TO HARVEST

Begin by cleaning empty bins thoroughly to ensure that the storage facility is free of leftover grains from previous year(s). Clean harvest equipment, augers and belts to eliminate potential infestations from old insect-infested grain. Using a low-pressure sprayer, apply your selected Bug Free Grains product as a surface, spot and crack-and-crevice treatment both inside and outside of bins (including any equipment).

### PRODUCT RECOMMENDATIONS INCLUDE:

| Diacon® IGR *PLUS* and PBO-8® Synergist or Gravista®-D Insecticide

### TREATING GRAINS DURING LOADING\*

Set up product application equipment prior to the loading of grains. As grain is loaded into facilities, calibrate equipment to allow a uniform distribution of grain protectants. Grains can be treated based on your storage strategy. It is imperative to knock down existing insects and protect against future generations.

\*Refer to product information and label for solutions that meet your insect control/storage strategy needs.



## TREATING EXISTING INFESTATIONS

If the infested grain is already in storage, fumigate with an EPA registered fumigant to eliminate the infestation. Following fumigation, if possible, turn the grain and treat to protect it from future infestations. Treatment recommendations will depend on your storage strategy.

## PRODUCT RECOMMENDATIONS INCLUDE:\*

**Short-term strategy:** Centynal™ Synergized Insecticide

**Long-term strategy:** Gravista® Insecticide or Gravista®-D Insecticide

\*Refer to product information and label for solutions that meet insect control/storage strategy needs.

## PROACTIVE INSECT MANAGEMENT

We have the most comprehensive mix of product solutions to meet your grain storage strategy needs. From pre-bin and rescue treatments to short- and long-term insect control needs, our formulations can help meet your goals and protect your bottom line.



## GRAIN TREATMENT TIPS & TRICKS

- Keep a daily record of applications to stored grain. Using either a digital or paper record, write down the date, amount of grain treated and amount of chemical used.
- Proper droplet size during product application will help eliminate drift. Droplets should be coarse or very coarse, measuring at least 157-360 microns.
- Select the correct nozzle type to control spray droplet size; a flat-fan or cone nozzle with a coarse spray will yield optimally sized droplets.
- Keep spray pressure at 20-25 psi to prevent droplets from drifting away from grains. Calibrate nozzle flow rates at the recommended pressure to help ensure proper grain coverage.
- When positioning your spraying equipment in relation to the grain stream, it should be sprayed in a manner that hits the grain and not the adjoining equipment. Test the direction and formation of your spray pattern right as you begin, and make adjustments as necessary. Adding more nozzles to improve coverage is a plus.
- Whether you're spraying a tank-mix of multiple products or a simple dilution, the differing weights of the liquids can cause separation or settling of active ingredients. That's why it's important to agitate your mixture frequently until spraying is complete.
- To help keep the mixture properly agitated, select a pump with enough volume to both apply and return the right amount of solution. Adding mechanical agitation to the tank is also a viable option.



# OUR PRODUCTS

The Bug Free Grains lineup of products has the most comprehensive selection of stored grain protectants in the industry. We continue to evolve our solutions to meet the needs of producers and stored grain professionals.



## INSECT GROWTH REGULATORS



Insect growth regulators (IGRs) disrupt the insect life cycle by interrupting larval development. Diacon® IGR products are tolerance exempt IGRs that can be used everywhere stored product insects are a problem. They can be used on all stored grains, spices, feed and seeds including those used for animal and human consumption. From commercial elevator and on-site farm storage to food processing facilities and everywhere in between, Diacon® IGR products go where insects are to provide **LONG-TERM CONTROL AND PROFIT PROTECTION.**



40 lb box

### Diacon®-D IGR

A dry formulation insect growth regulator designed for locations where water is scarce, Diacon®-D IGR is a ready-to-use option for controlling insects at on-farm and off-premise storage sites.

#### FEATURES AND BENEFITS

- Ready-to-use formulation
- Labeled for all stored commodities including malting barley
- EPA tolerance exempt to control insect larvae in food commodities
- Contains the active ingredient (S)-methoprene
- Interrupts listed insects' life cycles for long-term control
- Can be applied to the grain stream or as an empty premise treatment
- CODEX approved for cereal grains
- Targets almond moths, Indian meal moths, lesser grain borers, sawtoothed grain beetles, merchant grain beetles, red and confused flour beetles, and other listed insect pests



2.5 gal, 1 gal and 1 liter

### Diacon® IGR

Diacon® IGR is a versatile liquid formulation that is excellent for control of lesser grain borers, Indian meal moths, sawtoothed grain beetles, flour beetles and other listed insect pests.

#### FEATURES AND BENEFITS

- Broad-spectrum control
- EPA tolerance exempt to control insect larvae in food commodities
- Contains the active ingredient (S)-methoprene
- Can be applied directly to grain stream
- Can be applied as a fogging concentrate or a surface spray
- CODEX approved for cereal grains
- CODEX approved for peanuts
- Targets almond moths, Indian meal moths, lesser grain borers, sawtoothed grain beetles, merchant grain beetles, red flour beetles, confused flour beetles and other listed insect pests

**CODEX approval is significant to allow countries to import Diacon®-treated commodities; however, not all countries accept CODEX MRLs. Check with the country of export to determine if CODEX MRLs are accepted.**

# ADULTICIDES



Adulticides target adult insects as opposed to insects in their larval stages. Bug Free Grains Stored Product Solutions lineup of products offers adulticides with two types of active ingredients: pyrethrins naturally derived from *Chrysanthemum cinerariifolium* and deltamethrin. Both provide a quick kill with residual control.



1 gal

## Centynal™ EC Insecticide

Get efficacious control over stored product insects with Centynal™ EC Insecticide. Effective when used alone or tank-mixed with Diacon® IGR and/or PBO-8® Synergist, Centynal™ EC Insecticide can be applied directly to stored grains or as an empty-bin treatment as part of proper sanitation practices.

### FEATURES AND BENEFITS

- Provides effective knockdown and control of stored product insects
- Contains the active ingredient deltamethrin
- For use on wheat, corn, rice and other labeled grains
- Enhanced emulsifiable concentrate formulation
- May be diluted with water or oil
- Miscible in dust-controlling oils
- Low-toxicity profile
- May be tank-mixed with Diacon® IGR and PBO-8® Synergist
- Broad-use pattern
- CODEX approved for cereal grains
- Targets lesser grain borers, larger grain borers, Angoumois grain moths, Indian meal moths, tobacco moths, dermestids, bean weevils, granary weevils, rice weevils, maize weevils, sawtoothed grain beetles, red flour beetles and confused flour beetles



1 gal and 1 qt

## Pyronyl™ Crop Spray

Pyronyl™ Crop Spray is a concentrated pyrethrum insecticide that features a high ratio of active ingredient to synergist for the flushing and rapid knockdown of multiple stored product insects. Use it alone or tank-mixed with Diacon® IGR as an effective tool for applications in and around stored grain facilities.

### FEATURES AND BENEFITS

- Controls stored product insects
- Contains synergized pyrethrins
- Ideal for the rapid knockdown of multiple species
- Quick degradation in the environment
- May be used as a fogging agent in conjunction with Diacon® IGR
- Targets ants, cockroaches, cadelles, cigarette beetles, confused flour beetles, dark mealworms, dried fruit beetles, drugstore beetles, grain mites, red flour beetles, rice weevils, sawtoothed grain beetles, spider beetles, yellow mealworms, Angoumois grain moths, Indian meal moths, mosquitoes, Mediterranean flour moths, small flying moths, tobacco moths and other listed insect pests



# SYNERGISTS

While synergists can display limited pesticidal activity on their own, their value comes in enhancing the potency of other pesticides. Some synergists work by blocking the enzymes insects use to metabolize the active ingredients of certain insecticides.



1 gal

## PBO-8® Synergist

PBO-8® Synergist improves the efficacy of insecticides. Use it as the perfect tank-mix partner with Centynal™ EC Insecticide or Diacon® IGR PLUS to provide an extra boost for stronger stopping power.

### FEATURES AND BENEFITS

- Blocks insects' ability to metabolize adulticides
- Synergizes many insecticides, including pyrethroids
- Works with most misting, fogging or spray equipment
- Presents a low risk to humans and environment
- Is an excellent resistance management tool



# OUR PRODUCTS

## COMBINATION PRODUCTS

Central Life Sciences offers numerous combination products featuring grain protectant technology that helps control larval development as well as kills adult insects. Our combination products can add to the knockdown power of an adulticide with the boost of a synergist for challenging infestations and/or the long-term control of an insect growth regulator (IGR), all in a single product. These products are formulated to contain the right ratio of active ingredients to control insects and are a convenient alternative to tank-mixing.



2.5 gal

### Gravista<sup>®</sup> Insecticide

Featuring 3-in-1 insect control with an adulticide, an insect growth regulator (IGR) and a synergist, Gravista<sup>®</sup> Insecticide kills exposed insects and breaks the life cycles. Gravista<sup>®</sup> Insecticide can be diluted with water or oil and can be applied directly on the grain stream or as an empty-bin and warehouse treatment.

#### FEATURES AND BENEFITS

- Three active ingredients: (S)-methoprene, Deltamethrin, Piperonyl Butoxide
- Broad-spectrum and long-term residual control
- On-grain, empty-bin and perimeter\* treatment
- For use on wheat, corn, rice and other labeled grains and seeds
- Water, oil and dust-controlling oils dilutable formulation
- CODEX approved for cereal grains
- May be tank-mixed with other products such as Pyronyl™ Crop Spray
- Targets lesser grain borers, larger grain borers, Angoumois grain moths, Indian meal moths, tobacco moths, Dermestids, and bean weevils; also, to control populations of granary weevils, rice weevils, maize weevils, sawtoothed grain beetles, red flour beetles and confused flour beetles

\*Not for use in California.



44 lb box

### Gravista<sup>®</sup>-D Insecticide

Gravista<sup>®</sup>-D Insecticide combines a synergized adulticide with an insect growth regulator. This dry formulation product can be applied directly on post-harvest grain streams. Gravista<sup>®</sup>-D Insecticide is an ideal solution to protect on-farm stored grains from the elusive weevil and other labeled insects.

#### FEATURES AND BENEFITS

- Combines a synergized adulticide with an insect growth regulator
- Ready-to-use formulation
- Non-restricted use
- 11 lb per 1,000 bu of grain
- Helps control weevils and other labeled pests
- For use on grain sorghum (milo), corn, popcorn, barley, oats, rice, rye and wheat
- Application to the grain stream and/or as an empty bin treatment
- CODEX approved for cereal grains
- Targets Indian meal moths, sawtoothed grain beetles, red flour beetles, rice weevils, maize weevils, lesser grain borers and other listed insect pests



2.5 gal

## Centynal™ Synergized Insecticide

Centynal™ Synergized Insecticide brings a boost to short-term grain storage and rescue treatments. By combining both an adulticide and a synergist, Centynal™ Synergized Insecticide provides effective knockdown and will help control even the toughest stored product insects.

### FEATURES AND BENEFITS

- Insecticide plus synergist that increases speed and knockdown of most labeled insects
- Tough on hard-to-control stored product insects
- CODEX approved for cereal grains
- Reduces rebound of infestations
- For use on wheat, corn, rice and other labeled grains and seeds
- On-grain, empty-bin and perimeter treatment
- Targets lesser grain borers, larger grain borers, Angoumois grain moths, Indian meal moths, tobacco moths, Dermestids, bean weevils, granary weevils, rice weevils, maize weevils, sawtoothed grain beetles, red flour beetles and confused flour beetles



2.5 gal and 1 qt

## Diacon® IGR PLUS

Diacon® IGR PLUS provides control of stored product insects through all life stages. Diacon® IGR PLUS can be applied directly to stored grains and used for treating empty bins and warehouses for stored product insect control.

### FEATURES AND BENEFITS

- Helps eliminate labeled stored product insects
- For use on wheat, corn, rice and other labeled grains
- Ideal for both on-grain use and sanitation
- Active ingredients control both adult and larval stages of insects
- Miscible in dust-controlling oils
- May be diluted with water or oil
- May be tank-mixed with PBO-8® Synergist for weevil control
- Long-residual control
- Reduces rebound of infestations
- CODEX approved for listed commodities
- Targets lesser grain borers, larger grain borers, Angoumois grain moths, Indian meal moths, tobacco moths, Dermestids, bean weevils, granary weevils, rice weevils, maize weevils, sawtoothed grain beetles, red flour beetles and confused flour beetles

# PREMISE CONTROL



This all-in-one product is formulated to provide insect control all over your storage facility. Containing a long-lasting insect growth regulator (IGR) and a broad-spectrum insecticide with a synergist, this three-in-one product provides fogging and space spray flexibility, ideal for mills, food-handling establishments, warehouses and more.



1 gal

## Inselux® Fog & Mill Spray

Featuring an insect growth regulator, an adulticide and a synergist, Inselux® Fog & Mill Spray kills exposed labeled insects and breaks the insects' life cycles for long-term control. It may be used undiluted or diluted in equipment capable of applying a fine mist. The product may be applied through cold foggers, thermal foggers, backpack misters or other equipment capable of applying a space spray.

### FEATURES AND BENEFITS

- Three active ingredients; (S)-methoprene, Pyrethrins, Piperonyl Butoxide
- May be diluted or undiluted
- For use in grain, feed, food processing and storage areas
- IGR reduces rebound of infestations
- Broad-spectrum control
- See label for controlled listed insect pests

**CODEX approval is significant to allow countries to import Diacon®-treated commodities; however, not all countries accept CODEX MRLs. Check with the country of export to determine if CODEX MRLs are accepted.**



# APPLICATION & DILUTION



## APPLICATION RATES\*

Amount of Centynal™ EC Insecticide/1,000 Bushels of Grain				
Commodity	0.5 ppm	1 ppm	Gallons of Water	Food Grade Mineral Oil/Gallon
	fl. oz.	fl. oz.	gal.	gal.
Barley	7.75	15.5	3-5	Follow oil manufacturer's recommendation
Corn	9	18	3-5	
Oats	5.15	10.3	3-5	
Popcorn	9	18	3-5	
Rice	7.2	14.4	3-5	
Rye	9	18	3-5	
Sorghum (milo)	8	16	3-5	
Wheat	9.6	19.2	3-5	

## DILUTION RATES

Centynal™ EC Insecticide Dilution Rates	
Application	
On-grain application with water as carrier	3-5 gal. solution per 1,000 bu.
On-grain application with food grade oil as carrier	Follow oil manufacturer's recommendation
Rough rice spray	10 gal. liquid per 1,000 bu.

Centynal™ EC Insecticide Tank Mix Rates	
PBO-8® Synergist	1 part Centynal™ EC Insecticide + 1.5 parts PBO-8® Synergist
Diacon® IGR	4 fl. oz. in mix to do 5 gallons solution per 1,000 bu.

## APPLICATION RATES\*

Amount of Centynal™ Synergized Insecticide/1,000 Bushels of Grain				
Commodity	Treatment at 0.5 ppm Deltamethrin/ 14 ppm PBO		Gallons of Diluent	Liters of Diluent
	fl. oz.	mL	gal.	L.
Barley	31	916	3-5	11.3-19
Corn	35.6	1,053	3-5	11.3-19
Oats	20.38	603	3-5	11.3-19
Popcorn	35.6	1,053	3-5	11.3-19
Rice	28.5	843	10	11.3-19
Rye	35.6	1,053	3-5	11.3-19
Sorghum (milo)	31.7	937	3-5	11.3-19
Wheat	38	1,124	3-5	11.3-19

Amount of Centynal™ Synergized Insecticide/Ton of Grain				
Commodity	Treatment at 0.5 ppm Deltamethrin/ 14 ppm PBO		Ounces of Diluent	Milliliters of Diluent
	fl. oz.	mL	fl. oz.	mL
Barley, Corn, Oats, Popcorn, Rice, Rye, Sorghum (milo), Wheat	1.3	38	13-21	384-630

## DILUTION RATES

Centynal™ Synergized Insecticide Dilution Rates	
Application	
On-grain application with water or food grade oil	3-5 gal. solution per 1,000 bu.
Rough rice spray	10 gal. of water per 1,000 bu.
Grain bins/warehouse	1-6 fl. oz. + 1 gal. water per 1,000 sq. ft.



# RATES



## APPLICATION RATES\*

### Diacon®-D IGR Application Rates

Application	
On grain application	8-10 lbs. per 1,000 bu.
Mixing into animal feed	0.33 lbs. per ton of feed
Empty bin treatment	1.5 oz. per 1,000 sq. ft.



## APPLICATION RATES\*

### Diacon® IGR Application Rates (Per 1,000 Bushels) in Solution

Commodity	fl. oz.
Wheat	7
Corn	7
Sorghum (milo)	7
Barley	6
Rice	6
Oats	4
Peanuts**	4
Sunflower	4

## DILUTION RATES

### Diacon® IGR Dilution Rates

On-grain application with water as carrier	5 gal. of solution per 1,000 bu.
On-grain application with food grade oil as carrier***	1 gal. of solution per 1,000 bu.
Rice spray with carrier as water	10 gal. of solution per 1,000 bu.
Fogging - aerosol generators****	1/10 fl. oz. per 10,000 cu. ft.



## APPLICATION RATES\*

### Amount of Diacon® IGR PLUS/1,000 Bushels of Grain

Commodity	0.5 ppm Deltamethrin/ 1.25 ppm (S)-Methoprene	1 ppm Deltamethrin/ 2.5 ppm (S)-Methoprene	Gallons of Water	Food Grade Mineral Oil/Gallon
	fl. oz.	fl. oz.	gal.	gal.
Barley	7.75	15.5	5	1
Corn	9	18	5	1
Oats	5.15	10.3	5	1
Popcorn	9	18	5	1
Rice	7.2	14.4	5	1
Rye	9	18	5	1
Sorghum (milo)	8	16	5	1
Wheat	9.6	19.2	5	1

## DILUTION RATES

### Diacon® IGR PLUS Dilution Rates

Application	
On-grain application	3-5 gal. liquid per 1,000 bu.
Rough rice spray	10 gal. liquid per 1,000 bu.
Surface spray	1 fl. oz. Diacon® IGR PLUS + 1.5 fl. oz. PBO-8® Synergist + water to make 1 gal. of solution per 1,000 sq. ft.

\*See label for complete product directions. \*\*For peanuts, use water solution only. \*\*\*Apply at the oil manufacturer's label application rate. \*\*\*\*Follow equipment manufacturer instructions for amount of diluent.



# APPLICATION & DILUTION



## APPLICATION RATES\*

Amount of Gravista®/1,000 Bushels of Grain				
Commodity	Treatment at 0.5 ppm Deltamethrin/ 14 ppm PBO/1.2 ppm (S)-Methoprene		Gallons of Diluent	Liters of Diluent
	fl. oz.	mL	gal.	L.
Barley	31	916	3-5	11.3-19
Corn	35.6	1,053	3-5	11.3-19
Oats	20.38	603	3-5	11.3-19
Popcorn	35.6	1,053	3-5	11.3-19
Rice	28.5	843	10	11.3-19
Rye	35.6	1,053	3-5	11.3-19
Sorghum (milo)	31.7	937	3-5	11.3-19
Wheat	38	1,124	3-5	11.3-19

Amount of Gravista®/Ton of Grain				
Commodity	Treatment at 0.5 ppm Deltamethrin/ 14 ppm PBO/1.2 ppm (S)-Methoprene		Ounces of Diluent	Milliliters of Diluent
	fl. oz.	mL	fl. oz.	mL
Barley, Corn, Oats, Popcorn, Rice, Rye, Sorghum (milo), Wheat	1.3	38	13-21	384-630

## DILUTION RATES

Gravista® Dilution Rates	
Application	
On-grain application with water or food grade oil	3-5 gal. solution per 1,000 bu.
Rough rice spray	10 gal. of water per 1,000 bu.
Grain bins/warehouse	1-6 fl. oz. + 1 gal. water per 1,000 sq. ft.



## APPLICATION RATES\*

Gravista®-D Insecticide Application Rates	
Application	
On-grain application (Barley, Corn, Oats, Popcorn, Rice, Rye, Grain Sorghum [milo], Wheat)	11 lbs. per 1,000 bu.
Empty bin treatment	13 oz. per 1,000 sq. ft.

# RATES



## APPLICATION RATES\*

To kill exposed stages of the species listed below, use the indicated rate of Inselux® Fog & Mill Spray per 1,000 cu. ft.

Insect	Rate of Inselux® Fog & Mill Spray in fluid ounces:
<ul style="list-style-type: none"> <li>House Flies, Indian Meal Moths</li> <li>Sawtoothed Grain Beetles</li> <li>Mosquitoes, Rice Weevils</li> </ul>	1 fl. oz.
<ul style="list-style-type: none"> <li>Almond Moths, Angoumois Grain Moths, Boxelder Bugs, Bran Bugs</li> <li>Cadelles, Carpet Beetles</li> <li>Cheese Mites, Cheese Skippers</li> <li>Chocolate Moths, Cigarette Beetles</li> <li>Clover Mites, Crickets</li> <li>Confused Flour Beetles, Drugstore Beetles, Earwigs, Flour Beetles, Grain Beetles, Grain Mites, Granary Weevils, Foreign Grain Beetles</li> </ul>	<ul style="list-style-type: none"> <li>Fruit Flies, Gnats, Non-Biting Ham Mites</li> <li>Lesser Grain Borers, Mealworms</li> <li>Mediterranean Flour Moths</li> <li>Merchant Grain Beetles</li> <li>Nuisance Ants**</li> <li>Red Flour Beetles, Rice Flour Beetles</li> <li>Rusty Grain Beetles, Silverfish</li> <li>Small Flying Moths, Spider Beetles</li> <li>Tobacco Moths, Warehouse Beetles</li> <li>Spiders***</li> </ul>

## DILUTION RATES

### Inselux® Fog & Mill Spray Dilution Rates

May be applied undiluted or diluted with mineral oil at a 1:1 ratio through mechanical, thermal or compressed air fogging systems, as a fogging mist or space spray.



## APPLICATION RATES\*

### Amount of PBO-8® Synergist/1,000 Bushels of Grain

Commodity	Mixing with Centynal™ EC	Mixing with Diacon® IGR PLUS at
	Insecticide at 0.5 ppm	0.5 ppm Deltamethrin/1.25 ppm (S)-Methoprene
	fl. oz.	fl. oz.
Barley	11.625	11.625
Corn	13.5	13.5
Oats	7.725	7.725
Popcorn	13.5	13.5
Rice	10.8	10.8
Rye	13.5	13.5
Sorghum (milo)	12	12
Wheat	14.4	14.4

\* See label for complete product directions.

\*\* Except carpenter, fire, harvester and pharaoh ants.

\*\*\* Except black widows and brown recluse spiders.



# Centynal<sup>EC</sup><sup>TM</sup>

## INSECTICIDE

**PROTECTS STORED GRAINS AND SEEDS AGAINST DAMAGE FROM INDIAN MEAL MOTH, SAW-TOOTHED GRAIN BEETLE, RED FLOUR BEETLE, RICE WEEVIL, MAIZE WEEVIL, AND MANY OTHER LISTED PESTS**

- Emulsifiable concentrate
- Residual control of listed pests
- May be diluted with water or oil
- Miscible in dust controlling oils
- May be tank-mixed

## SPECIMEN LABEL

### ACTIVE INGREDIENT:

Deltamethrin\* ..... 4.75%

**OTHER INGREDIENTS:** ..... 95.25%  
100.00%

\*Contains 0.41 lb deltamethrin per gallon

EPA Reg. No. 89459-87      EPA Est. No. 2724-TX-1

**KEEP OUT OF REACH OF CHILDREN  
WARNING**

See additional Precautionary Statements

### PRECAUTIONARY STATEMENTS – HAZARDS TO HUMANS & DOMESTIC ANIMALS – WARNING

May be fatal if swallowed. Causes substantial but temporary eye injury. Do not get in eyes, on skin or on clothing. Wear protective eyewear (goggles, face shield or safety glasses). Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove and wash contaminated clothing before reuse. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals.

### FIRST AID

Call a poison control center or doctor immediately for treatment advice.

- |                     |  |
|---------------------|--|
| <b>If swallowed</b> | <ul style="list-style-type: none"> <li>• Have person sip a glass of water if able to swallow.</li> <li>• Do not induce vomiting unless told to by a poison control center or doctor.</li> <li>• Do not give anything by mouth to an unconscious person.</li> </ul> |
|---------------------|--|

*(continued)*

<b>If in eyes</b>	<ul style="list-style-type: none"> <li>• Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li> <li>• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b>If on skin</b>	<ul style="list-style-type: none"> <li>• Take off contaminated clothing.</li> <li>• Rinse skin immediately with plenty of water for 15-20 minutes.</li> </ul>
<p>Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-248-7763 for emergency medical treatment information.</p>	

### ENVIRONMENTAL HAZARDS

This product is extremely toxic to freshwater and estuarine fish and invertebrates. To protect the environment, do not allow pesticide to enter or runoff into storm drains, drainage ditches, gutters, or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Rinsing application equipment over the treated area will help avoid runoff to water bodies or drainage systems.

This pesticide is highly toxic to bees exposed to direct treatment. Do not apply or allow it to drift to crops or weeds on which bees are actively foraging.

### PHYSICAL AND CHEMICAL HAZARDS

Do not use water-based sprays of **Centynal<sup>TM</sup> EC Insecticide (Centynal<sup>TM</sup> EC)** in conduits, motor housings, junction boxes, switch boxes, or other electrical equipment because of possible shock hazard.

### DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

## PRODUCT INFORMATION

**Centynal™ EC** is an emulsifiable concentrate insecticide for use both indoors and outdoors for effective knockdown and residual control of listed pests. Also for use on stored grains and for treating empty bins and warehouses for stored product pest control. **Centynal™ EC** provides effective knockdown and control of listed adults and larvae.

**Mixing Instructions:** Mix **Centynal™ EC** with water and apply using a hand pressurized or power operated sprayer. For outdoor broadcast application, apply as a fine or coarse spray using suitable equipment.

### Dilution Instructions:

- Add half the water to the spray tank
- Add **Centynal™ EC** as specified in the appropriate application rates table
- Agitate well to insure proper mixing
- Add the remaining water slowly

**Note:** Re-agitation of spray dilution may be required if left overnight. It is recommended that water be used as the diluent for applications in non-commodity treatments. Use a mixed dilution within 48 hours of preparing.

**Tank Mixing:** **Centynal™ EC** may be tank-mixed with other currently registered pesticides unless prohibited by the tank-mix partner label. To check for product compatibility, a small volume mixing test with other products is recommended. The more restrictive label will apply when tank-mixing.

### STORED GRAIN

For use to control adult and accessible stages of larvae of listed insect pests including, lesser grain borers, larger grain borers, Angoumois grain moths, Indian meal moths, tobacco moths, dermestids, and bean weevils; also to suppress populations of granary weevils, rice weevils, maize weevils, saw-toothed grain beetles, red flour beetles, and confused flour beetles.

In instances of high weevil complex pressure or suspected tolerance, the use of piperonyl butoxide (PBO) synergist may be used as a tank mix partner. Use PBO at a ratio of 1 part **Centynal™ EC** to 1.5 parts PBO. **Centynal™ EC** may be tank-mixed with other pesticide products such as Pyronyl™ Crop spray. The more stringent label will apply when tank-mixing

**Directions for Treatment:** Dilute **Centynal™ EC** with water or approved dust controlling oils, mineral or soybean oils and apply to the moving grain stream as a coarse spray to provide a concentration of 0.5 ppm - 1 ppm of **Centynal™ EC** on the grain.

For applications with water, use 3 – 5 gallons of liquid per 1,000 bushels of grain. Use more or less dilution volume depending on the situation but ensure that uniform coverage is obtained. For rough rice, use 10 gallons of water per 1,000 bushels. Maintain agitation before and during application. Use a mixed dilution within 48 hours of preparing.

Re-agitation of spray dilution may be required if left overnight.

When using oils as a diluent, follow the oil manufacturers' recommendations. For best results, apply the solution as uniformly as possible as the grain is placed into storage.

For stored grain, apply **Centynal™ EC** in solution at the labeled rate of 3 - 5 gallons of dilution per 1,000 bushels to achieve a 0.5 – 1 ppm concentration of **Centynal™ EC**.

Apply at rates as shown in the tables below:

AMOUNT OF CENTYNAL™ EC/1000 BUSHELS OF GRAIN						
Commodity	Centynal™ EC Treatment at 0.5 ppm		Centynal™ EC Treatment at 1 ppm		Gallons of Diluent	Liters of Diluent
	fl oz	ml	fl oz	ml		
Barley	7.75	229	15.50	458	3 – 5	11.3 - 19
Corn	9	266	18	532	3 – 5	11.3 – 19
Oats	5.15	152	10.30	304	3 – 5	11.3 – 19
Popcorn	9	266	18	532	3 – 5	11.3 - 19
Rice	7.20	212	14.40	424	3 – 5	11.3 – 19
Rye	9	266	18	532	3 – 5	11.3 – 19
Sorghum (milo)	8	236	16	472	3 – 5	11.3 – 19
Wheat	9.6	283	19.20	566	3 – 5	11.3 – 19

AMOUNT OF CENTYNAL™ EC/TON OF GRAIN						
Commodity	Centynal™ EC @ 0.5 ppm		Centynal™ EC @ 1 ppm		Ounces of Diluent	Milliliters of Diluent
	fl oz	ml	fl oz	ml		
Barley	0.34	10	0.68	20	13 – 21	384 – 630
Corn	0.34	10	0.68	20	13 – 21	384 – 630
Oats	0.34	10	0.68	20	13 – 21	384 – 630
Popcorn	0.34	10	0.68	20	13 – 21	384 – 630
Rice	0.34	10	0.68	20	13 – 21	384 – 630
Rye	0.34	10	0.68	20	13 – 21	384 – 630
Sorghum (milo)	0.34	10	0.68	20	13 – 21	384 – 630
Wheat	0.34	10	0.68	20	13 - 21	384 – 630

### SEED TREATMENT

Apply **Centynal™ EC** to protect stored grain seed against injury from stored grain insects listed. Clean and treat seed storage areas prior to storage. If seed is infested, fumigate\*\* to control existing populations and apply **Centynal™ EC** for residual protection. **Centynal™ EC** may also be used in conjunction with registered fungicides or other insecticide seed protectants, to protect barley, corn, oats, popcorn, rice, rye, sorghum and wheat seeds during the seed treating process, and prior to bagging or placing in bulk seed storage at the rate of 0.5 – 1 ppm. When treating seed with other pesticide products, properly tag the treated seed indicating: "Do not use the seed for food, feed, or oil purposes." Seed treated with **Centynal™ EC** alone is considered grain.

\*\*Fumigate with EPA-registered fumigant according to manufacturer's label recommendations.

### GRAIN BINS AND WAREHOUSES

Dilute 0.25 – 1.5 fl oz of **Centynal™ EC** with water to make 1 gallon of solution. Before storing or handling grain, apply finished spray to equipment, wall and floor surfaces of grain bins and warehouses at the rate of 1 gallon per 1,000 sq ft but do not allow runoff to occur. Apply after bins and buildings have been thoroughly cleaned. If infestations are severe or there are areas that are difficult to treat with a conventional spray, fumigation is recommended with an EPA-registered fumigant.

Use **Centynal™ EC** as a perimeter treatment to prevent infestation of these structures. Apply to areas where listed pests are seen or may find entrance. Treat a band of soil or turf up to 6-10 feet wide around and adjacent to structures and treat the foundation to a height of 2 to 3 feet. Mix 0.25 – 1.5 fl oz of **Centynal™ EC** with water to make 1 gallon of solution. Apply as a coarse spray at a rate of 1 gallon per 1,000 sq ft. Thoroughly and uniformly wet the barrier area but do not allow runoff to occur.

## PEST CONTROL OUTDOORS

Apply **Centynal™ EC** as a perimeter treatment, residual spray or broadcast application to lawns, grounds and ornamental plantings. Apply as a perimeter treatment or residual spray around non-residential settings. Begin treatments at the first sighting of listed pests.

**Residual Spray:** Apply **Centynal™ EC** as a coarse low-pressure spray diluted in water at the rate of 0.75 to 1.5 fl oz to surfaces of buildings, porches, screens, window frames, eaves, patios, refuse dumps, garages and lawns.

**Perimeter Treatments:** For best results, apply **Centynal™ EC** to areas where listed pests are seen and may find entrance. Treat a band of soil or turf up to 6-10 feet wide around and adjacent to buildings and treat the building foundation to a height of 2 to 3 feet. Apply as a coarse spray at the rate of 0.25 to 1.5 fl oz of **Centynal™ EC** in sufficient amount of water. (See Mixing Directions for Perimeter Application below). Use an adequate volume to thoroughly and uniformly wet the barrier area but do not allow runoff to occur.

### MIXING DIRECTIONS FOR CENTYNAL™ EC OUTDOOR PERIMETER APPLICATION

#### CENTYNAL™ EC DILUTION (0.25 fl oz/1000 sq ft)

Tank Size	15 Gallons		25 Gallons		50 Gallons		100 Gallons	
	Centynal™ EC Per Tank (fl oz)	Structures Per Tank*	Centynal™ EC Per Tank (fl oz)	Structures Per Tank*	Centynal™ EC Per Tank (fl oz)	Structures Per Tank*	Centynal™ EC Per Tank (fl oz)	Structures Per Tank*
1	3.75	15	6.25	25	12.50	50	25.00	100
3	1.25	5	2.08	8	4.17	16	8.33	33
5	0.75	3	1.25	5	2.50	10	5.00	20
10	0.37	1.5	0.63	2.5	1.25	5	2.50	10

\*Based on **Centynal™ EC** applied at 0.25 fl oz per 1,000 sq ft per structure

#### CENTYNAL™ EC DILUTION (1 fl oz/1000 sq ft)

Tank Size	15 Gallons		25 Gallons		50 Gallons		100 Gallons	
	Centynal™ EC Per Tank (fl oz)	Structures Per Tank*	Centynal™ EC Per Tank (fl oz)	Structures Per Tank*	Centynal™ EC Per Tank (fl oz)	Structures Per Tank*	Centynal™ EC Per Tank (fl oz)	Structures Per Tank*
1	15.0	15	25.0	25	50.0	50.0	100.0	100.0
3	5.0	5	8.4	8.3	16.7	16.7	33.3	33.3
5	3.0	3	5.0	5	10.0	10.0	20.0	20.0
10	1.5	1.5	2.5	2.5	5.0	5	10.0	10.0

\*Based on **Centynal™ EC** applied at 1 fl oz per 1,000 sq ft per structure

Do not apply **Centynal™ EC** until the location of heating or air conditioning ducts, vents, water and sewer lines and electrical conduits are identified. Once known, applications may be made around these structures, however, be careful to avoid contamination of these structures and airways. Do not apply through any type of irrigation system. Do not water the treated area to the point of runoff. Do not make applications during rain.

**STORAGE and DISPOSAL:** Do not contaminate water, food, or feed by storage or disposal. **PESTICIDE STORAGE** – Mix as needed. Store in original container in a secured, dry storage area. Avoid extremes in temperatures. Prevent cross-contamination with other pesticides and fertilizers. If the container is leaking and/or material is spilled, on floor or paved surfaces, absorb on sawdust or other commercially available absorbing material, sweep up and remove to chemical waste area for your disposal. Concentrate is stable at normal storage temperatures. **PESTICIDE DISPOSAL** - Wastes resulting from use of this product may be disposed of on site or at an approved waste disposal facility. **CONTAINER HANDLING** – Nonrefillable container. Do not reuse or refill this container. Triple rinse (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

**IMPORTANT: READ BEFORE USE**

Read the entire Directions for Use, Conditions, Disclaimer of Warranties and Limitations of Liability before using this product. If terms are not acceptable, return the unopened product container at once. By using this product, user or buyer accepts the following Conditions, Disclaimer of Warranties and Limitations of Liability.

**CONDITIONS:** The Directions for Use of this product are believed to be adequate and must be followed carefully. However, because of manner of use and other factors beyond Central Garden & Pet Company's control, it is impossible for Central Garden & Pet Company to eliminate all risks associated with the use of this product. As a result, crop injury or ineffectiveness is always possible. All such risks shall be assumed by the user or buyer.

**DISCLAIMER OF WARRANTIES:** To the extent consistent with applicable law, CENTRAL GARDEN & PET COMPANY MAKES NO OTHER WARRANTIES, EXPRESS OR IMPLIED, OR MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE OR OTHERWISE, THAT EXTEND BEYOND THE STATEMENTS MADE ON THIS LABEL. No agent of Central Garden & Pet Company is authorized to make any warranties beyond those contained herein or to modify the warranties container herein. To the extent consistent with applicable law, Central Garden & Pet Company disclaims any liability whatsoever for special, incidental or consequential damages resulting from the use or handling of this product.

**LIMITATIONS OF LIABILITY:** TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER IN CONTRACT, WARRANTY, TORT, NEGLIGENCE, STRICT LIABILITY OR OTHERWISE, SHALL NOT EXCEED THE PURCHASE PRICE PAID, OR AT CENTRAL GARDEN & PET COMPANY'S ELECTION, THE REPLACEMENT OF PRODUCT.

For information or in case of an emergency, call 1-800-248-7763.

[www.bugfreegrains.com](http://www.bugfreegrains.com)



## Combination synergist and adulticide

**PROTECTS STORED GRAINS, SEEDS, GRAIN BINS AND WAREHOUSES (INDOOR AND PERIMETER\* TREATMENTS) AGAINST DAMAGE FROM INDIAN MEAL MOTH, SAWTOOTHED GRAIN BEETLE, RED FLOUR BEETLE, RICE WEEVIL, MAIZE WEEVIL, AND OTHER LISTED PESTS**

- Emulsifiable concentrate
- For use on grains, empty bin and perimeter\* treatment
- Residual control
- May be diluted with water or oil
- Miscible in dust-controlling oils

### ACTIVE INGREDIENT:

Deltamethrin .....	1.2%
Piperonyl butoxide .....	33.3%
<b>OTHER INGREDIENTS:</b> .....	<b>65.5%</b>
<b>TOTAL</b>	<b>100.0%</b>

Contains 0.1 lb deltamethrin per gallon  
 Contains 2.67 lb piperonyl butoxide per gallon  
 \*Not for use in California

DELTAMETHRIN	GROUP	3A	INSECTICIDE
--------------	-------	----	-------------

# SPECIMEN LABEL

EPA Reg. No. 89459-115

EPA Est. No. 2724-TX-1

## KEEP OUT OF REACH OF CHILDREN WARNING

See additional Precautionary Statements

### PRECAUTIONARY STATEMENTS – HAZARDS TO HUMANS & DOMESTIC ANIMALS – WARNING

May be fatal if swallowed. Causes moderate eye irritation. Avoid contact with eyes, skin, or clothing. Wear protective eyewear (goggles, face shield or safety glasses). Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet. Remove and wash contaminated clothing before reuse.

#### Personal Protective Equipment (PPE):

Mixers, loaders, applicators and other handlers must wear:  
 Long-sleeved shirt, long pants,  
 Shoes and socks

Chemical-resistant gloves: barrier laminate, nitrile rubber ≥ 14 mils, neoprene rubber ≥ 14 mils, or Viton ≥ 14 mils

In addition to the above PPE, applicators using a high pressure handwand in an enclosed area must wear a minimum of a NIOSH-approved particulate filtering facepiece respirator with any R or P filter; OR a NIOSH-approved elastomeric particulate respirator with any R or P filter; OR a NIOSH-approved powered air purifying respirator with HE filters.

**User Safety Requirements:** Follow manufacturer’s instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry. Discard clothing and other absorbent materials that have been drenched or heavily contaminated with the product’s concentrate. Do not reuse them.

**User Safety Recommendations:** Users should wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Users should remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. Users should remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

### FIRST AID

<b>If swallowed</b>	<ul style="list-style-type: none"> <li>• Call a poison control center or doctor immediately for treatment advice.</li> <li>• Have person sip a glass of water if able to swallow.</li> <li>• Do not induce vomiting unless told to by a poison control center or doctor.</li> <li>• Do not give anything by mouth to an unconscious person.</li> </ul>
---------------------	--

<b>If on skin</b>	<ul style="list-style-type: none"> <li>• Take off contaminated clothing.</li> <li>• Rinse skin immediately with plenty of water for 15-20 minutes.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b>If in eyes</b>	<ul style="list-style-type: none"> <li>• Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li> <li>• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<p>Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-800-248-7763 for emergency medical treatment information.</p>	

### ENVIRONMENTAL HAZARDS

This product is extremely toxic to aquatic organisms, including fish and invertebrates. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters, or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Rinsing application equipment over the treated area will help avoid run off to water bodies or drainage systems.

This pesticide is highly toxic to bees exposed to direct treatment on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds while bees are foraging the treatment area.

### PHYSICAL AND CHEMICAL HAZARDS

Do not use water-based sprays of **Centynal™ Synergized Insecticide (Centynal™ Synergized)** in conduits, motor housings, junction boxes, switch boxes, or other electrical equipment because of possible shock hazard.

### DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

### RESISTANCE-MANAGEMENT

For resistance management, please note that **Centynal™ Synergized** contains a Group 3A insecticide. Any insect population may contain individuals naturally resistant to **Centynal™ Synergized** and other Group 3A insecticides. The resistant individuals may dominate the insect population if these insecticides are used repeatedly in the same fields. Appropriate resistance-management strategies should be followed. To delay insecticide/acaricide resistance, take the following steps:

- Use tank mixtures with insecticides from a different group that are equally effective on the target pest when such use is permitted. Do not rely on the same mixture repeatedly for the same pest population. Consider any known cross-resistance issues (for the targeted pests) between the individual components of a mixture.

In addition, consider the following recommendations provided by the Insecticide Resistance Action Committee (IRAC):

- Individual insecticides selected for use in mixtures should be highly effective and be applied at the rates at which they are individually registered for use against the target species.
- Mixtures with components having the same IRAC mode of action classification are not recommended for insect resistance management. When using mixtures, consider any known cross-resistance issues between the individual components for the targeted pest(s).
- Mixtures become less effective if resistance is already developing to one or both active ingredients, but they may still provide pest management benefits.
- The insect resistance management benefits of an insecticide mixture are greatest if the two components have similar periods of residual insecticidal activity. Mixtures of insecticides with unequal periods of residual insecticide activity may offer an insect resistance management benefit only for the period where both insecticides are active.
- Monitor after application for unexpected target pest survival. If the level of survival suggests the presence of resistance, consult with your local university specialist or certified pest control advisor.
- Contact your local extension specialist for any additional pesticide resistance-management and/or IPM recommendations for the specific site and pest problems in your area.
- For further information or to report suspected resistance contact Central Garden & Pet Company at 1-800-248-7763 or at [www.bugfreegrains.com](http://www.bugfreegrains.com).

### PRODUCT INFORMATION

**Centynal™ Synergized** is an emulsifiable concentrate insecticide that contains an adulticide and synergist for use on stored grains and for treating empty bins and warehouses for stored product pest control. **Centynal™ Synergized** provides effective knockdown and control of listed adults. For both indoor and outdoor use.

### USE RESTRICTIONS

Do not remain in treated area. Exit area immediately and remain outside the treated area until aerosols, vapors, and/or mists have dispersed.

Do not enter or allow others to enter until sprays have dried. Do not apply this product in a way that will contact workers or other persons, children or pets, either directly or through drift. Only protected handlers may be in the area during applications. Do not allow adults, children, or pets to enter the treated area until sprays have dried.

Do not allow adults, children, or pets to enter until vapors, mists and aerosols have dispersed and the treated area has been thoroughly ventilated.

Do not apply more than one time per day. Remove or cover exposed food and drinking water before application.

Remove or cover dishes, utensils, food processing equipment and food preparation surfaces, or wash them before use.

Do not use in aircraft cabins except in compliance with PR Notice 96-3.

When used in dairy barns or facilities, close milk bulk tank lids to prevent contamination from spray and from dead or falling insects. Remove or cover milking utensils before application.

Wash teats of animals before milking.

Do not apply directly into sewers or drains or to any area like a gutter where drainage to storm sewers, water bodies, or aquatic habitat can occur.

Do not allow the product to enter any drain during or after application.

Not for use in outdoor residential misting systems.

All outdoor spray applications must be limited to spot or crack-and-crevice treatments only, except for the following permitted uses

1. Application to pervious surfaces such as soil, lawn, turf, and other vegetation;
2. Perimeter band treatments of 7 feet wide or less from the base of a man-made structure to pervious surfaces (e.g., soil, mulch, or lawn);
3. Applications to underside of eaves, soffits, doors, or windows permanently protected from rainfall by a covering, overhang, awning, or other structure;
4. Applications around potential exterior pest entry points into man-made structures such as doorways and windows, when limited to a band not to exceed one inch;
5. Applications to vertical surfaces (such as the side of a man-made structure) directly above impervious surfaces (e.g., driveways, sidewalks, etc.), up to 2 feet above ground level;
6. Applications to vertical surfaces directly above pervious surfaces, such as soil, lawn, turf, mulch or other vegetation) only if the pervious surface does not drain into ditches, storm drains, gutters, or surface waters.

Spot treatments must not exceed two square feet in size (for example, 2 ft. by 1 ft. or 4 ft. by 0.5 ft.).

For soil or foliar applications, do not apply by ground within 25 feet of lakes, reservoirs, rivers, permanent streams, marshes or natural ponds, estuaries and commercial fish farm ponds.

Do not spray the product into fish pools, ponds, streams, or lakes. Do not apply directly to sewers or storm drains, or to any area like a drain or gutter where drainage to sewers, storm drains, water bodies, or aquatic habitat can occur.

Do not allow the product to enter any drain during or after application.

Do not apply directly to impervious horizontal surfaces such as sidewalks, driveways, and patios except as a spot or crack-and-crevice treatment.

Treat surfaces to ensure thorough coverage but avoid runoff.

To treat insects harbored in voids and cracks-and-crevices, applications must be made in such a manner to limit dripping and avoid runoff onto untreated structural surfaces and plants.

Do not apply or irrigate to the point of runoff. Do not make applications during rain. Avoid making applications when rainfall is expected before the product has sufficient time to dry

(minimum 4 hours). Rainfall within 24 hours after application may cause unintended runoff of pesticide application.

### STORED GRAIN

Use to control adults and accessible larval stages of listed insect pests including, lesser grain borers, larger grain borers, Angoumois grain moths, Indian meal moths, tobacco moths, Dermestids, and bean weevils; also, to control populations of granary weevils, rice weevils, maize weevils, sawtoothed grain beetles, red flour beetles, and confused flour beetles.

#### Tank-mixing

**Centynal™ Synergized** may be tank-mixed with other pesticide products such as Pyronyl™ Crop Spray and or Diacon® IGR. It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and Directions for Use on all product labels involved in tank-mixing. Users must follow the most restrictive Directions for Use and Precautionary Statements of each product in the tank mixture.

#### Directions for Treatment

Dilute **Centynal™ Synergized** with water or approved dust-controlling oils, mineral or soybean oils and apply to the moving grain stream as a coarse spray.

Final spray volume with water solutions is 3 – 5 gallons of liquid per 1,000 bushels of grain. For rough rice, use 10 gallons of water per 1,000 bushels. Maintain agitation before and during application. Use a mixed dilution within 48 hours of preparing. Re-agitation of spray dilution may be required if left overnight.

For final spray volumes with oil solutions, follow the oil manufacturer's rate recommendations.

For uniform application, treat the grain as it moves into storage. Apply **Centynal™ Synergized** in solution at the rate of 3 – 5 gallons of dilution per 1,000 bushels to achieve a 0.5 ppm on the commodity. Apply at rates as shown in the following tables:

#### AMOUNT OF CENTYNAL™ SYNERGIZED/ 1,000 BUSHELS OF GRAIN

Commodity	TREATMENT AT 0.5 ppm Deltamethrin/14 ppm PBO		Gallons of Diluent	Liters of Diluent
	fl oz	mL		
Barley	31	916	3 – 5	11.3 – 19
Corn	35.6	1053	3 – 5	11.3 – 19
Oats	20.38	603	3 – 5	11.3 – 19
Popcorn	35.6	1053	3 – 5	11.3 – 19
Rice	28.5	843	10	11.3 – 19
Rye	35.6	1053	3 – 5	11.3 – 19
Sorghum (milo)	31.7	937	3 – 5	11.3 – 19
Wheat	38	1124	3 – 5	11.3 – 19

## AMOUNT OF CENTYNAL™ SYNERGIZED/ TON OF GRAIN

Commodity	TREATMENT AT 0.5 ppm Deltamethrin/14 ppm PBO		Ounces of Diluent	Milliliters of Diluent
	fl oz	mL	fl oz	mL
Barley	1.3	38	13 – 21	384 – 630
Corn	1.3	38	13 – 21	384 – 630
Oats	1.3	38	13 – 21	384 – 630
Popcorn	1.3	38	13 – 21	384 – 630
Rice	1.3	38	13 – 21	384 – 630
Rye	1.3	38	13 – 21	384 – 630
Sorghum (milo)	1.3	38	13 – 21	384 – 630
Wheat	1.3	38	13 – 21	384 – 630

### SEED TREATMENT

Apply **Centynal™ Synergized** to protect stored grain seed against injury from listed insect pests. Clean and treat seed storage areas prior to storage. If seed is infested, fumigate\*\* to control existing populations and apply **Centynal™ Synergized** for residual protection. **Centynal™ Synergized** may also be used in conjunction with registered fungicides or other insecticide seed protectants, to protect barley, corn, oats, popcorn, rice, rye, sorghum and wheat seed during the seed treating process, and prior to bagging or placing in bulk seed storage. When treating seed with other pesticide products, properly tag the treated seed, indicating "Do not use seed for food, feed, or oil purposes." Seed treated with **Centynal™ Synergized** alone is considered grain.

\*\*Fumigate with EPA-registered fumigants according to manufacturer's label recommendations.

### GRAIN BINS AND WAREHOUSES

Dilute 1–6 fl oz of **Centynal™ Synergized** with water to make 1 gallon of solution. Before storing or handling grain, apply finished spray to equipment, wall and floor surfaces of grain bins and warehouses at the rate of 1 gallon per 1,000 sq ft but do not allow runoff to occur. Apply after equipment bins and buildings have been thoroughly cleaned. If infestations are severe or there are areas that are difficult to treat with a conventional spray, fumigation is recommended with an EPA-registered fumigant.

Use **Centynal™ Synergized** as a perimeter\* treatment to prevent infestation of these structures. Apply to areas where listed pests are seen or may find entrance. Treat a band of soil or grass up to 6–7 feet wide around and adjacent to structures and treat the foundation to a height of 2 feet. Mix 1–6 fl oz of **Centynal™ Synergized** with water to make 1 gallon of solution. Apply as a coarse spray at a rate of 1 gallon per 1,000 sq ft. Wet the barrier area thoroughly and uniformly but do not allow runoff to occur. All outdoor\* applications to impervious surfaces are limited to spot and crack-and-crevice applications only.

\*Not for use in California

**Central Garden & Pet Company**  
1501 East Woodfield Road 200W  
Schaumburg, Illinois 60173

AG 21-023

**STORAGE AND DISPOSAL:** Do not contaminate water, food, or feed by storage or disposal. **PESTICIDE STORAGE** – Mix as needed. Store in original container in a secured, dry storage area. Avoid extremes in temperatures. Prevent cross-contamination with other pesticides and fertilizers. If the container is leaking and/or material is spilled, on floor or paved surfaces, absorb on sawdust or other commercially available absorbing material, sweep up and remove to chemical waste area for your disposal. Concentrate is stable at normal storage temperatures. **PESTICIDE DISPOSAL** – Wastes resulting from use of this product may be disposed of on site or at an approved waste disposal facility. **CONTAINER HANDLING** – Nonrefillable container. Do not reuse or refill this container. Triple rinse (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or reconditioning if appropriate, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

### IMPORTANT: READ BEFORE USE

Read the entire Directions for Use, Conditions, Disclaimer of Warranties and Limitations of Liability before using this product. If terms are not acceptable, return the unopened product container at once. By using this product, user or buyer accepts the following Conditions, Disclaimer of Warranties and Limitations of Liability.

**CONDITIONS:** The Directions for Use of this product are believed to be adequate and must be followed carefully. However, because of manner of use and other factors beyond Central Garden & Pet Company's control, it is impossible for Central Garden & Pet Company to eliminate all risks associated with the use of this product. As a result, crop injury or ineffectiveness is always possible. All such risks shall be assumed by the user or buyer.

**DISCLAIMER OF WARRANTIES:** To the extent consistent with applicable law, CENTRAL GARDEN & PET COMPANY MAKES NO OTHER WARRANTIES, EXPRESS OR IMPLIED, OR MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE OR OTHERWISE, THAT EXTEND BEYOND THE STATEMENTS MADE ON THIS LABEL. No agent of Central Garden & Pet Company is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. To the extent consistent with applicable law, Central Garden & Pet Company disclaims any liability whatsoever for special, incidental or consequential damages resulting from the use or handling of this product.

**LIMITATIONS OF LIABILITY:** TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER IN CONTRACT, WARRANTY, TORT, NEGLIGENCE, STRICT LIABILITY OR OTHERWISE, SHALL NOT EXCEED THE PURCHASE PRICE PAID, OR AT CENTRAL GARDEN & PET COMPANY'S ELECTION, THE REPLACEMENT OF PRODUCT.

For information or in case of an emergency, call 1-800-248-7763. Visit our Web site: [www.bugfreegrains.com](http://www.bugfreegrains.com)

Centynal, Diacon and Pyronyl are trademarks of Wellmark International.  
©2021 Wellmark International.

September, 2021  
Schaumburg, IL

# Diacon-D

IGR

- **AN INSECT GROWTH REGULATOR TO CONTROL STORED PRODUCT INSECTS BY BREAKING THE PEST LIFE CYCLE AND PREVENTING LARVAE FROM MATURING INTO ADULTS**
- APPROVED FOR USE ON ALL GRAINS, SPICES, FEEDS, AND SEEDS, INCLUDING COMMODITIES FOR HUMAN AND ANIMAL CONSUMPTION
- VERSATILE APPLICATION INCLUDES – DIRECT TO GRAIN, TOP-DRESS, MIXING INTO ANIMAL FEEDS, OR EMPTY PREMISE TREATMENTS

## SPECIMEN LABEL

### ACTIVE INGREDIENT:

(S)-Methoprene (CAS# 65733-16-6)..... 0.8%

**OTHER INGREDIENTS:**\* .....99.2%

TOTAL ..... 100.0%

EPA Reg. No. 2724-788

EPA Est. No. 2724-TX-1

**KEEP OUT OF REACH OF CHILDREN**

**CAUTION**

See additional Precautions

### PRECAUTIONARY STATEMENTS – HAZARDS TO HUMANS & DOMESTIC ANIMALS CAUTION

Harmful if inhaled or absorbed through the skin. Causes moderate eye irritation. Avoid breathing dust. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet. Remove contaminated clothing and wash before reuse. Wear chemical resistant gloves, protective eyewear, and dust mask.

**Personal Protective Equipment (PPE):** Handlers and applicators must wear a dust mask, chemical-resistant gloves, and protective eyewear.

### FIRST AID

**IF INHALED** • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. • Call a poison control center or doctor for treatment advice.

**IF ON SKIN** • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.

**IF IN EYES** • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eyes. • Call a poison control center or doctor for treatment advice.

**IF SWALLOWED** • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Do not give anything to an unconscious person.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-248-7763 for emergency medical treatment information.

### ENVIRONMENTAL HAZARDS

Do not apply directly to water or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater or rinsate.

### GENERAL INFORMATION

**DIACON®-D IGR** is a ready-to-use formulation that can either be applied to commodities and feed prior to storage, top-dressed to commodities already stored, or used as an empty premise treatment for areas where commodities will be stored. The residual activity of **DIACON®-D IGR** prevents the emergence of stored product insects during storage of commodities/feed. (S)-Methoprene, the active ingredient in **DIACON®-D IGR** protects the stored commodity from damaging insects by interfering with the normal process of insect development. Unlike traditional pesticides, **DIACON®-D IGR** is not an adulticide, but its residual activity prevents the development of larval insects into viable reproducing adults. **DIACON®-D IGR**, when used according to the label directions, prevents adult emergence and subsequent new generations of insects. **DIACON®-D IGR** is effective against

insect pests, such as: almond moth, Indian meal moth, lesser grain borer, sawtooth grain beetle, merchant grain beetle, red and confused flour beetle, and other insect pests that are affected by Insect Growth Regulators (IGR's). If the commodity is known to be infested when it is initially being placed into storage, fumigate the commodity to eliminate the existing insect populations following filling of bins or other storage facilities. Then apply **DIACON®-D IGR** for residual protection.

Apply **DIACON®-D IGR** to any food commodity (including but not limited to: cereal grains, corn, sunflower, canola, legumes, popcorn, wheat, spices, sorghum, rice, cocoa, peanuts, oats, millet and processed foods readied for packing) for control of insect larvae. **DIACON®-D IGR** may also be used to treat bulk or bagged pet food, livestock feed, birdseed, or any other animal feed. **DIACON®-D IGR** may be applied to any seed stock. Treated commodities may be processed immediately. **Applications of DIACON®-D IGR to finished animal feeds, birdseed or other finished products DO NOT require package labeling since the application is made to control pests in the package and no other claims are made on the final package.**

### DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

### DIRECTIONS FOR TREATMENT OF COMMODITIES PRIOR TO STORAGE

For protection of stored food, grain, animal feed, seeds used for oil, and seed stock against stored product insects - for optimum results, thoroughly clean and treat storage areas prior to storing commodities. **DIACON®-D IGR** may be used in conjunction with aeration.

Wearing a dust mask and protective gloves, apply **DIACON®-D IGR** to grain or other commodities as they are being loaded or turned into storage facilities. Apply 8-10 pounds of **DIACON®-D IGR** per 1000 bushels of commodity. Apply the dust as uniformly as possible to the commodity stream to assure even coverage. Apply in such a manner to reduce the potential of the product blowing off target in windy conditions. The higher application rate will provide longer residual.

### DIRECTIONS FOR TOP-DRESS TREATMENT OF STORED GRAIN

Use **DIACON®-D IGR** as a top-dressing application for commodities which were previously stored in bins or other storage areas. To apply as a top-dressing; wearing a dust mask and protective gloves, uniformly apply the **DIACON®-D IGR** over the commodity's surface. Dust the bin headspace paying particular attention to the top of the commodity mass, beams, ceilings, and rafters. Use equipment capable of applying a uniform layer of **DIACON®-D IGR**. Thorough coverage is essential.

As a top-dressing application, apply 8 pounds of **DIACON®-D IGR** per 1000 ft<sup>2</sup> of commodity surface area. Raking the top layer will enhance protection from invading insects infesting the top surfaces of the commodity. When raking commodity surface, rake to a depth of 1 foot.

### DIRECTIONS FOR MIXING INTO ANIMAL FEEDS

**DIACON®-D IGR** is suitable to be used in the production of bulk or bagged animal feeds (including but not limited to: pet food, livestock feed, birdseed, and all other animal feeds). Wearing a dust mask and protective gloves, add **DIACON®-D IGR** to achieve a thorough mix within the feed. Apply 0.33 pounds (150 gms) – 0.26 pounds (126 gms) per ton of animal feed and mix to ensure uniformity. Treated feeds may be fed immediately with no withdrawal required.

### DIRECTIONS FOR EMPTY BIN, WALL VOID, CRACK AND CREVICE TREATMENT

Use **DIACON®-D IGR** to treat bins, warehouses, mills, food processing and packaging facilities, feed mills, and other areas where stored product pests may harborage. For applications to hard surfaces within these sites, wearing a dust mask and protective gloves, apply **DIACON®-D IGR** at a rate of 1.5 oz (45 grams) per 1000 ft<sup>2</sup> of surface area. Pay particular attention to cracks, crevices, and voids where food commodities can accumulate and attract pests. Use a bulb duster or other suitable equipment to apply the material.

### STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal. **Storage** - Store in a cool, dry place. **Pesticide Disposal** - Wastes resulting from use of this product may be disposed of on site or at an approved waste disposal facility. **Container Disposal** – Nonrefillable container. Do not reuse or refill this container. Completely empty bag into application equipment. Then dispose of empty bag and container in a sanitary landfill, or by incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

### WARRANTY AND CONDITIONS OF SALE

To the fullest extent permitted by law, Seller makes no warranty, expressed or implied, concerning the use and handling of this product other than indicated on the label. Buyer assumes all risks of use and handling of this material when such use and handling are contrary to label instructions.

For information or in case of an emergency, call 1-800-248-7763.

[www.bugfreegrains.com](http://www.bugfreegrains.com)

**Wellmark International**  
1501 East Woodfield Road 200W  
Schaumburg, Illinois 60173

Diacon and Diacon with design are trademarks of Wellmark International.  
©2012 Wellmark International.

April, 2012  
Schaumburg, IL



- AN INSECT GROWTH REGULATOR TO CONTROL INSECTS IN STORED FOOD COMMODITIES
- APPROVED FOR USE ON ALL GRAINS, SPICES, FEEDS, AND SEEDS FOR HUMAN AND ANIMAL CONSUMPTION

# SPECIMEN LABEL

### ACTIVE INGREDIENT:

(S)-Methoprene (CAS# 65733-16-6)..... 33.6%

**OTHER INGREDIENTS:**\* ..... 66.4%

TOTAL ..... 100.0%

\*Contains petroleum distillates

This product contains 2.5 lb/gallon (288 grams/liter) (S)-Methoprene active ingredient

EPA Reg. No. 2724-427

EPA Est. No. 2724-TX-1

**KEEP OUT OF REACH OF CHILDREN**

**CAUTION**

See additional Precautions

### PRECAUTIONARY STATEMENTS – HAZARDS TO HUMANS & DOMESTIC ANIMALS CAUTION

Causes moderate eye irritation. Harmful if absorbed through skin. Avoid contact with eyes, skin, or clothing. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash clothing before reuse.

#### FIRST AID

**IF IN EYES** • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eyes. • Call a poison control center or doctor for treatment advice.

**IF ON SKIN** • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.

**IF SWALLOWED** • Immediately call a poison control center or doctor. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give ANY liquid to the person. Do not give anything by mouth to an unconscious person.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact Wellmark at 1-800-248-7763 (24 hr/7 days) for emergency medical treatment information.

**NOTE TO PHYSICIAN:** Contains petroleum distillates – vomiting may cause aspiration pneumonia.

### GENERAL INFORMATION

**GENERAL INFORMATION:** (S)-Methoprene, the active ingredient in DIACON® IGR, protects stored grains, agricultural commodities, seed, and animal feeds from damaging insects by interfering with the normal process of insect development. Unlike traditional pesticides, DIACON® IGR is not an adulticide, but its residual activity prevents the development of larvae into adults. When used according to the label directions, DIACON® IGR inhibits growth of larvae of these insects: almond moth, Indian meal moth, lesser grain borer, sawtoothed grain beetle, merchant grain beetle, red flour beetle, confused flour beetle, and other insect pests that are affected by Insect Growth Regulators (IGR's). Treat existing insect populations with an adulticide, then apply DIACON® IGR for residual protection.

### DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Apply DIACON® IGR to any food commodity (including but not limited to: cereal grains, corn, sunflower, canola, legumes, popcorn, wheat, spices, sorghum, rice, cocoa, peanuts, oats, millet, and processed or non-processed commodities readied for packaging) for control of insect larvae. DIACON® IGR may also be used to treat bulk or bagged pet food, livestock feed components and finished livestock feed, birdseed, or any other dry animal feed. DIACON® IGR may be applied to any seed stock. Treated commodities may be processed immediately.

## NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Do not remain in treated areas and ventilate before re-entry. Leave the room closed for 30 minutes to allow spray or fog mist to settle before re-entry into the area.

### DIRECTIONS FOR TREATMENT OF GRAIN BEING STORED:

For protection of stored food, grain, animal feed, seeds used for oil, and seed stock against insects - for optimum results, thoroughly clean and treat storage areas prior to storing commodities.

**DILUTION INSTRUCTIONS:** DIACON® IGR may be diluted in water, or FDA-approved food grade oils, or soybean oil. Final water or oil dilution volume is 5 gallons of liquid per 1,000 bushels of grain and 10 gallons of liquid per 1,000 bushels of rice (or 21 fl oz per ton for grain, or 42 fl oz per ton for rice). Apply diluted product to the moving grain stream as a uniform coarse spray. Lower or greater volumes of diluent can be used but, it is important to assure that the commodity mass is uniformly treated. When using oil dilutions, apply at the oil manufacturer's label application rates. Agitate water or oil solutions to ensure thorough mixing.

Apply DIACON® IGR in solution in accordance with the general directions, to stored products using the following tables as a guide. A range of rates for different commodities is provided for flexibility to address varying storage periods and conditions. DIACON® IGR may be applied at any rate within the specified range indicated on the tables. However, for maximum residual activity, use higher rates; the lower application rates provide shorter residual activity.

#### APPLICATION RATES (PER 1,000 BUSHELS)

COMMODITY	fl oz	ml	fl oz	ml	fl oz	ml
Wheat	7	210	3.5	105	1.75	52
Corn	7	210	3.5	105	1.75	52
Sorghum (Milo)	7	210	3.5	105	1.75	52
Barley	6	180	3.0	90	1.50	45
Rice	6	180	3.0	90	1.50	45
Oats	4	120	2.0	60	1.00	30
Peanuts**	4	120	2.0	60	1.00	30
Sunflower	4	120	2.0	60	1.00	30

\*\*For peanuts, use water solution only

#### APPLICATION RATES (PER TON OF PRODUCT)

COMMODITY	fl oz	ml	fl oz	ml	fl oz	ml
Any	0.25	7.5	0.125	3.7	0.0625	1.8

To determine application rates for other commodities, contact your distributor or Central Life Sciences.

### USE NOTES

- Apply DIACON® IGR grain protectant only to grain of known treatment history.
- Check spray equipment calibration regularly to ensure proper application rates.

- Add the measured amount of DIACON® IGR to a spray tank filled with ½ gallon of water or oil, mix, and complete filling with specified quantity of diluent. Keep mixed solution out of direct sunlight.
- Agitate before each use.
- Pre-treat storage facilities before filling with DIACON® IGR treated commodities.
- DIACON® IGR may be used in conjunction with aeration.
- DIACON® IGR may be used in metering systems in conjunction with dust-controlling oils. Follow the equipment and oil manufacturers' directions; ensure that the label rate of DIACON® IGR is applied uniformly.
- Applications of DIACON® IGR to finished animal feeds, birdseed, or other finished products DO NOT require package labeling since the application is made to control pests in the package and no other claims are made on the final packaging.
- DIACON® IGR may be tank-mixed with a suitable adulticide in diluted solutions or when fogging. When tank-mixing DIACON® IGR with an adulticide, the more stringent label will apply.
- DIACON® IGR can be used with dust-controlling oils or finishing oils used in bird feed.

### DIRECTIONS FOR TOP-DRESS TREATMENT OF STORED GRAIN:

Apply DIACON® IGR as a top-dressing to stored products placed in bins or storage areas. For top-dressing application, spray or fog the bin headspace paying particular attention to the top of the commodity mass, beams, ceilings, and rafters. Thorough coverage of the bins or storage areas is essential for optimal results. For surface treatments, apply 1 ml (1/30 fl oz) of DIACON® IGR per 1,000 square feet; or for space spray, apply 3 ml (1/10 fl oz) per 10,000 cubic feet in sufficient diluent to provide adequate coverage. Do not flood top-dress area.

### DIRECTIONS FOR MIXING INTO BULK OR BAGGED ANIMAL FEEDS:

DIACON® IGR is suitable to be used in the production of bulk or bagged animal feeds (including but not limited to: pet food, livestock feed, birdseed, and other animal feeds). Add 3.7-15 ml (0.125 to 0.5 fl oz) of DIACON® IGR per ton of animal feed with a sufficient diluent and mix to achieve thorough coverage.

### DIRECTIONS FOR FOGGING OR SPRAYING:

DIACON® IGR may also be used as a fogging concentrate or surface spray for the treatment of insect pests that are affected by Insect Growth Regulators: infesting warehouses, silos, storage bins, or other stored commodity areas. Apply DIACON® IGR in food processing, food service, food preparation, and food handling establishments including, mills, bakeries, restaurants, taverns, food storage facilities, meat and produce canneries, cafeterias, breweries, candy processing, pet food production, grocery stores, rail cars, ships, trucks, and trailers, cereal processing and bottling facilities. In addition to treating raw commodities and their processing facilities, apply DIACON® IGR to packaged, canned, and bottled foodstuffs, as well as the packaging such as boxes, liners, and pallets. For application of DIACON® IGR as a fogging or space spray, use the following directions.

**Application with Aerosol Generators:** Apply DIACON® IGR as an ultra low volume (ULV) mist or aerosol. Dilute with water or oil as required following the instructions of the spray equipment manufacturer. Apply DIACON® IGR at the rate of 1 ml (1/30 fl oz) per 1,000 square feet of surface area or 3 ml (1/10 fl oz) per 10,000 cubic feet. Use appropriate spray equipment, such as mechanical break up cold aerosol generators or thermal foggers, to achieve uniform coverage.

Close room or warehouse and shut off all ventilating systems. Apply the product as a fine mist in the air above the trays, shelves, and in areas where dust may accumulate. Do not remain in treated areas and ventilate before re-entry. Leave the room closed for 30 minutes to allow spray mist to settle before re-entry. Repeat application as required to suppress pest populations.

**Application as an Aqueous Spray:** To prepare a diluted spray solution, fill the mixing container with ½ gallon water, add 1 ml (1/30 fl oz) of DIACON® IGR, mix, and complete filling with a total of 1 gallon of water. Apply at a rate of 1 gallon of diluted solution per 1,000 square feet of surface area with any low-pressure sprayer typically used for indoor applications. Treat all areas that may harbor insect pests. Agitate solution before each use. Repeat application as required to suppress pest populations.

## STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

**Pesticide Storage** - Store in a cool, dry place.

**Pesticide Disposal** - Wastes resulting from use of this product may be disposed of on site or at an approved waste disposal facility.

**Container Disposal** - Nonrefillable container. Do not reuse or refill this container. Clean container promptly after emptying. Triple rinse as follows. Empty the remaining contents into application equipment or a mix tank and drain for ten seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for ten seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for ten seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available, or puncture and dispose of in a sanitary landfill, or by incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

## WARRANTY AND CONDITIONS OF SALE

To the extent consistent with applicable law, seller makes no warranty concerning the use and handling of this product other than indicated on the label. To the extent consistent with applicable law, buyer assumes all risks of use and handling of this material when such use and handling are contrary to label instructions.

For product information, call 1-800-248-7763,  
M - F, 8 am - 5 pm CST.

[www.bugfreegrains.com](http://www.bugfreegrains.com)



# Diacon<sup>®</sup> IGR PLUS

Combination insect growth regulator **PLUS** adulticide

**PROTECTS STORED GRAINS AND SEEDS AGAINST DAMAGE FROM INDIAN MEAL MOTH, SAW-TOOTHED GRAIN BEETLE, RED FLOUR BEETLE, RICE WEEVIL, MAIZE WEEVIL, AND OTHER LISTED PESTS**

- Long residual control
- Reduces rebound of infestations
- May be tank-mixed
- May be diluted with water or oil
- Miscible in dust controlling oils

## SPECIMEN LABEL

### ACTIVE INGREDIENT:

(Deltamethrin ..... 4.75%  
S-Methoprene ..... 11.40%

**OTHER INGREDIENTS:** ..... 83.85%

**TOTAL** ..... 100.00%

Contains 0.41 lb deltamethrin per gallon  
Contains 1 lb S-methoprene per gallon

EPA Reg. No. 89459-86

EPA Est. No. 2724-TX-1

**KEEP OUT OF REACH OF CHILDREN**

**WARNING**

See additional Precautions

### PRECAUTIONARY STATEMENTS – HAZARDS TO HUMANS & DOMESTIC ANIMALS – WARNING

May be fatal if swallowed. Causes substantial but temporary eye injury. Do not get in eyes, on skin, or on clothing. Wear protective eyewear (goggles, face shield or safety glasses). Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet. Remove and wash contaminated clothing before reuse. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals.

#### FIRST AID

<b>If swallowed</b>	<ul style="list-style-type: none"> <li>• Call a poison control center or doctor immediately for treatment advice.</li> <li>• Have person sip a glass of water if able to swallow.</li> <li>• Do not induce vomiting unless told to by a poison control center or doctor.</li> <li>• Do not give anything to an unconscious person.</li> </ul>
<b>If in eyes</b>	<ul style="list-style-type: none"> <li>• Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li> <li>• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>

(continued)

<b>If on skin</b>	<ul style="list-style-type: none"> <li>• Take off contaminated clothing.</li> <li>• Rinse skin immediately with plenty of water for 15-20 minutes.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<p>Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-248-7763 for emergency medical treatment information.</p>	

### ENVIRONMENTAL HAZARDS

This product is extremely toxic to freshwater and estuarine fish and invertebrates. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters, or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Rinsing application equipment over the treated area will help avoid run off to water bodies or drainage systems.

This pesticide is highly toxic to bees exposed to direct treatment. Do not apply or allow it to drift to crops or weeds on which bees are actively foraging.

### PHYSICAL AND CHEMICAL HAZARDS

Do not use water-based sprays of **DIACON<sup>®</sup> IGR PLUS** in conduits, motor housings, junction boxes, switch boxes, or other electrical equipment because of possible shock hazard.

### DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

### PRODUCT INFORMATION

**DIACON<sup>®</sup> IGR PLUS** is an emulsifiable concentrate insecticide that contains an adulticide and insect growth regulator for use on stored grains and for treating empty bins and warehouses for stored product pest control. **DIACON<sup>®</sup> IGR PLUS** provides effective knockdown and control of listed adults and larvae

and prevents the buildup of listed pests with long residual control provided by the IGR.

### STORED GRAIN

Use to control adults and accessible larval stages of listed insect pests including, lesser grain borers, larger grain borers, Angoumois grain moths, Indian meal moths, tobacco moths, dermestids, and bean weevils; also to control populations of granary weevils, rice weevils, maize weevils, saw-toothed grain beetles, red flour beetles, and confused flour beetles.

**Tank-mixing:** In instances of high weevil complex pressure or suspected tolerance, the use of piperonyl butoxide (PBO) synergist may be used as a tank mix partner. Use PBO at a ratio of 1 part **DIACON® IGR PLUS** to 1.5 parts PBO. **DIACON® IGR PLUS** may be tank-mixed with other pesticide products such as Pyronyl™ Crop Spray. The more stringent label will apply when tank-mixing.

**Directions for Treatment:** Dilute **DIACON® IGR PLUS** with water or approved dust-controlling oils, mineral or soybean oils and apply to the moving grain stream as a coarse spray to provide a concentration of 0.5 ppm to 1 ppm of deltamethrin and 1.25 ppm or 2.5 ppm of S-methoprene on the grain.

Final spray volume with water solutions is 3–5 gallons of liquid per 1,000 bushels of grain. For rough rice, use 10 gallons of water per 1,000 bushels. Maintain agitation before and during application. Use a mixed dilution within 48 hours of preparing. Re-agitation of spray dilution may be required if left overnight.

For final spray volumes with oil solutions, follow the oil manufacturer's rate recommendations.

For uniform application, treat the grain as it moves into storage. Apply **DIACON® IGR PLUS** in solution at the rate of 3–5 gallons of dilution per 1,000 bushels to achieve a 0.5 ppm to 1 ppm concentration of deltamethrin and 1.25 ppm or 2.5 ppm of S-methoprene on the commodity. Apply at rates as shown in the tables below:

### AMOUNT OF DIACON® IGR PLUS/ 1,000 BUSHELS OF GRAIN

Commodity	Treatment at 0.5 ppm Deltamethrin/ 1.25 ppm S-Methoprene		Treatment at 1 ppm Deltamethrin/ 2.5 ppm S-Methoprene		Gallons of Diluent	Liters of Diluent
	fl oz	ml	fl oz	ml		
Barley	7.75	229	15.50	458	3 – 5	11.3 – 19
Corn	9	266	18	532	3 – 5	11.3 – 19
Oats	5.15	152	10.30	304	3 – 5	11.3 – 19
Popcorn	9	266	18	532	3 – 5	11.3 – 19
Rice	7.2	212	14.40	424	3 – 5	11.3 – 19
Rye	9	266	18	532	3 – 5	11.3 – 19
Sorghum (milo)	8	236	16	472	3 – 5	11.3 – 19
Wheat	9.6	283	19.20	566	3 – 5	11.3 – 19

### AMOUNT OF DIACON® IGR PLUS /TON OF GRAIN

Commodity	0.5 ppm Deltamethrin/ 1.25 ppm S-Methoprene		1 ppm Deltamethrin/ 2.5 ppm S-Methoprene		Ounces of Diluent	Milliliters of Diluent
	fl oz	ml	fl oz	ml		
Barley	0.34	10	0.68	20	13 – 21	384 – 630
Corn	0.34	10	0.68	20	13 – 21	384 – 630
Oats	0.34	10	0.68	20	13 – 21	384 – 630
Popcorn	0.34	10	0.68	20	13 – 21	384 – 630
Rice	0.34	10	0.68	20	13 – 21	384 – 630
Rye	0.34	10	0.68	20	13 – 21	384 – 630
Sorghum (milo)	0.34	10	0.68	20	13 – 21	384 – 630
Wheat	0.34	10	0.68	20	13 – 21	384 – 630

### SEED TREATMENT

Apply **DIACON® IGR PLUS** to protect stored grain seed against injury from stored grain insects listed. Clean and treat seed storage areas prior to storage. If seed is infested, fumigate\*\* to control existing populations and apply **DIACON® IGR PLUS** for residual protection. **DIACON® IGR PLUS** may also be used in conjunction with registered fungicide or other insecticide seed protectants, to protect barley, corn, oats, popcorn, rice, rye, sorghum and wheat seed during the seed treating process, and prior to bagging or placing in bulk seed storage at the rate of 0.5 ppm – 1 ppm. When treating seed with other pesticide products, properly tag the treated seed, indicating "Do not use seed for food, feed, or oil purposes." Seed treated with **DIACON® IGR PLUS** alone is considered grain.

\*\*Fumigate with EPA-registered fumigants according to manufacturer's label recommendations.

### GRAIN BINS AND WAREHOUSES

Dilute 0.25 – 1.5 fl oz of **DIACON® IGR PLUS** with water to make 1 gallon of solution. Before storing or handling grain, apply finished spray to equipment, wall and floor surfaces of grain bins and warehouses at the rate of 1 gallon per 1,000 sq ft but do not allow runoff to occur. Apply after equipment bins and buildings have been thoroughly cleaned. If infestations are severe or there are areas that are difficult to treat with a conventional spray, fumigation is recommended with an EPA-registered fumigant.

Use **DIACON® IGR PLUS** as a perimeter treatment to prevent infestation of these structures. Apply to areas where listed pests are seen or may find entrance. Treat a band of soil or lawn up to 6–10 feet wide around and adjacent to structures and treat the foundation to a height of 2 to 3 feet. Mix 0.25–1.5 fl oz of **DIACON® IGR PLUS** with water to make 1 gallon of solution. Apply as a coarse spray at a rate of 1 gallon per 1,000 sq ft. Wet the barrier area thoroughly and uniformly but do not allow runoff to occur. Applications to building structures higher than 3 feet; all outdoor applications to impervious surfaces are limited to spot and crack-and-crevice applications only.

## STORAGE and DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

**PESTICIDE STORAGE** – Mix as needed. Store in original container in a secured, dry storage area. Avoid extremes in temperatures. Prevent cross-contamination with other pesticides and fertilizers. If the container is leaking and/or material is spilled, on floor or paved surfaces, absorb on sawdust or other commercially available absorbing material, sweep up and remove to chemical waste area for your disposal. Concentrate is stable at normal storage temperatures.

**PESTICIDE DISPOSAL** - Wastes resulting from use of this product may be disposed of on site or at an approved waste disposal facility.

**CONTAINER DISPOSAL** – Nonrefillable container. Do not reuse or refill this container. Triple rinse (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

## IMPORTANT: READ BEFORE USE

Read the entire Directions for Use, Conditions, Disclaimer of Warranties and Limitations of Liability before using this product. If terms are not acceptable, return the unopened product container at once. By using this product, user or buyer accepts the following Conditions, Disclaimer of Warranties and Limitations of Liability.

**CONDITIONS:** The Directions for Use of this product are believed to be adequate and must be followed carefully. However, because of manner of use and other factors beyond Central Garden & Pet Company's control, it is impossible for Central Garden & Pet Company to eliminate all risks associated with the use of this product. As a result, crop injury or ineffectiveness is always possible. All such risks shall be assumed by the user or buyer.

**DISCLAIMER OF WARRANTIES:** To the extent consistent with applicable law, CENTRAL GARDEN & PET COMPANY MAKES NO OTHER WARRANTIES, EXPRESS OR IMPLIED, OR MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE OR OTHERWISE, THAT EXTEND BEYOND THE STATEMENTS MADE ON THIS LABEL. No agent of Central Garden & Pet Company is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. To the extent consistent with applicable law, Central Garden & Pet Company disclaims any liability whatsoever for special, incidental or consequential damages resulting from the use or handling of this product.

**LIMITATION OF LIABILITY:** TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER IN CONTRACT, WARRANTY, TORT, NEGLIGENCE, STRICT LIABILITY OR OTHERWISE, SHALL NOT EXCEED THE PURCHASE PRICE PAID, OR AT CENTRAL GARDEN & PET COMPANY'S ELECTION, THE REPLACEMENT OF PRODUCT.

For information or in case of an emergency, call 1-800-248-7763.

[www.bugfreegrains.com](http://www.bugfreegrains.com)

**Central Garden & Pet Company**  
1501 East Woodfield Road 200W  
Schaumburg, Illinois 60173

Diacon and Diacon with design are trademarks of Wellmark International.  
©2016 Wellmark International.

November, 2016  
Schaumburg, IL



**Combination insect growth regulator, synergist and adulticide**

**PROTECTS STORED GRAINS, SEEDS, GRAIN BINS AND WAREHOUSES (INDOOR AND PERIMETER\* TREATMENTS) AGAINST DAMAGE FROM INDIAN MEAL MOTH, SAWTOOTHED GRAIN BEETLE, RED FLOUR BEETLE, RICE WEEVIL, MAIZE WEEVIL, AND OTHER LISTED PESTS**

- Three-way insect control
- For use on grains, empty bin and perimeter\* treatment
- Long residual control
- Kills listed adult insects and exposed larvae
- May be diluted with water or oil
- Miscible in dust-controlling oils

**ACTIVE INGREDIENTS:**

Deltamethrin .....	1.20%
Piperonyl butoxide .....	33.30%
S-Methoprene .....	2.85%
<b>OTHER INGREDIENTS:</b> .....	62.65%
	100.00%

Contains 0.1 lb deltamethrin per gallon. Contains 0.229 lb S-Methoprene per gallon.  
 Contains 2.67 lb piperonyl butoxide per gallon  
 \*Not for use in California

DELTAMETHRIN	GROUP	3A	INSECTICIDE
--------------	-------	----	-------------

**SPECIMEN LABEL**

EPA Reg. No. 89459-116 EPA Est. No. 2724-TX-1

**KEEP OUT OF REACH OF CHILDREN  
 WARNING**

See additional Precautionary Statements

**PRECAUTIONARY STATEMENTS – HAZARDS TO HUMANS & DOMESTIC ANIMALS – WARNING**

May be fatal if swallowed. Causes moderate eye irritation. Avoid contact with eyes, skin, or clothing. Wear protective eyewear (goggles, face shield or safety glasses). Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet. Remove and wash contaminated clothing before reuse.

**Personal Protective Equipment (PPE):**

Mixers, loaders, applicators and other handlers must wear:  
 Long-sleeved shirt, long pants,  
 Shoes and socks  
 Chemical-resistant gloves: barrier laminate, nitrile rubber ≥ 14 mils, neoprene rubber ≥ 14 mils, or Viton ≥ 14 mils  
 In addition to the above PPE, applicators using a high pressure handwand in an enclosed area must wear a minimum of a NIOSH-approved particulate filtering facepiece respirator with any R or P filter; OR a NIOSH-approved elastomeric particulate respirator with any R or P filter; OR a NIOSH-approved powered air purifying respirator with HE filters.

**User Safety Requirements:** Follow manufacturer’s instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry. Discard clothing and other absorbent materials that have been drenched or heavily contaminated with the product’s concentrate. Do not reuse them.

**User Safety Recommendations:** Users should wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Users should remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. Users should remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

**FIRST AID**

<b>If swallowed</b>	<ul style="list-style-type: none"> <li>• Call a poison control center or doctor immediately for treatment advice.</li> <li>• Have person sip a glass of water if able to swallow.</li> <li>• Do not induce vomiting unless told to by a poison control center or doctor.</li> <li>• Do not give anything by mouth to an unconscious person.</li> </ul>
<b>If on skin</b>	<ul style="list-style-type: none"> <li>• Take off contaminated clothing.</li> <li>• Rinse skin immediately with plenty of water for 15–20 minutes.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>

<b>If in eyes</b>	<ul style="list-style-type: none"> <li>• Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li> <li>• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
-------------------	---

Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-800-248-7763 for emergency medical treatment information.

### ENVIRONMENTAL HAZARDS

This product is extremely toxic to aquatic organisms, including fish and invertebrates. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters, or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Rinsing application equipment over the treated area will help avoid run off to water bodies or drainage systems.

This pesticide is highly toxic to bees exposed to direct treatment on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds while bees are foraging the treatment area.

### PHYSICAL AND CHEMICAL HAZARDS

Do not use water-based sprays of **Gravista® Insecticide (Gravista®)** in conduits, motor housings, junction boxes, switch boxes, or other electrical equipment because of possible shock hazard.

### DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

### RESISTANCE-MANAGEMENT

For resistance management, please note that **Gravista®** contains a Group 3A insecticide. Any insect population may contain individuals naturally resistant to **Gravista®** and other Group 3A insecticides.

The resistant individuals may dominate the insect population if these insecticides are used repeatedly in the same fields. Appropriate resistance-management strategies should be followed. To delay insecticide/acaricide resistance, take the following steps:

- Use tank mixtures with insecticides from a different group that are equally effective on the target pest when such use is permitted. Do not rely on the same mixture repeatedly for the same pest population. Consider any known cross-resistance issues (for the targeted pests) between the individual components of a mixture.

In addition, consider the following recommendations provided by the Insecticide Resistance Action Committee (IRAC):

- Individual insecticides selected for use in mixtures should be highly effective and be applied at the rates at which they are individually registered for use against the target species.

- Mixtures with components having the same IRAC mode of action classification are not recommended for insect resistance management. When using mixtures, consider any known cross-resistance issues between the individual components for the targeted pest(s).
- Mixtures become less effective if resistance is already developing to one or both active ingredients, but they may still provide pest management benefits.
- The insect resistance management benefits of an insecticide mixture are greatest if the two components have similar periods of residual insecticidal activity. Mixtures of insecticides with unequal periods of residual insecticide activity may offer an insect resistance management benefit only for the period where both insecticides are active.
- Monitor after application for unexpected target pest survival. If the level of survival suggests the presence of resistance, consult with your local university specialist or certified pest control advisor.
- Contact your local extension specialist for any additional pesticide resistance-management and/or IPM recommendations for the specific site and pest problems in your area.
- For further information or to report suspected resistance contact Central Garden & Pet Company at 1-800-248-7763 or at [www.bugfreegrains.com](http://www.bugfreegrains.com).

### PRODUCT INFORMATION

**Gravista®** is an emulsifiable concentrate insecticide that contains an adulticide, synergist and insect growth regulator for use on stored grains and for treating empty bins and warehouses for stored product pest control. **Gravista®** provides effective knockdown and control of listed adults and larvae and prevents the buildup of listed pests with long residual control provided by the IGR. For both indoor and outdoor use.

### USE RESTRICTIONS

Do not remain in treated area. Exit area immediately and remain outside the treated area until aerosols, vapors, and/or mists have dispersed.

Do not enter or allow others to enter until sprays have dried.

Do not apply this product in a way that will contact workers or other persons, children or pets, either directly or through drift. Only protected handlers may be in the area during applications.

Do not allow adults, children, or pets to enter the treated area until sprays have dried.

Do not allow adults, children, or pets to enter until vapors, mists and aerosols have dispersed and the treated area has been thoroughly ventilated.

Do not apply more than one time per day.

Remove or cover exposed food and drinking water before application.

Remove or cover dishes, utensils, food processing equipment and food preparation surfaces, or wash them before use.

Do not use in aircraft cabins except in compliance with PR Notice 96-3.

When used in dairy barns or facilities, close milk bulk tank lids to prevent contamination from spray and from dead or falling insects. Remove or cover milking utensils before application. Wash teats of animals before milking.

Do not apply directly into sewers or drains or to any area like a gutter where drainage to storm sewers, water bodies, or aquatic habitat can occur.

Do not allow the product to enter any drain during or after application.

Not for use in outdoor residential misting systems.

All outdoor spray applications must be limited to spot or crack-and-crevice treatments only, except for the following permitted uses

1. Application to pervious surfaces such as soil, lawn, turf, and other vegetation;
2. Perimeter band treatments of 7 feet wide or less from the base of a man-made structure to pervious surfaces (e.g., soil, mulch, or lawn);
3. Applications to underside of eaves, soffits, doors, or windows permanently protected from rainfall by a covering, overhang, awning, or other structure;
4. Applications around potential exterior pest entry points into man-made structures such as doorways and windows, when limited to a band not to exceed one inch;
5. Applications to vertical surfaces (such as the side of a man-made structure) directly above impervious surfaces (e.g., driveways, sidewalks, etc.), up to 2 feet above ground level;
6. Applications to vertical surfaces directly above pervious surfaces, such as soil, lawn, turf, mulch or other vegetation) only if the pervious surface does not drain into ditches, storm drains, gutters, or surface waters.

Spot treatments must not exceed two square feet in size (for example, 2 ft. by 1 ft. or 4 ft. by 0.5 ft.).

For soil or foliar applications, do not apply by ground within 25 feet of lakes, reservoirs, rivers, permanent streams, marshes or natural ponds, estuaries and commercial fish farm ponds.

Do not spray the product into fish pools, ponds, streams, or lakes. Do not apply directly to sewers or storm drains, or to any area like a drain or gutter where drainage to sewers, storm drains, water bodies, or aquatic habitat can occur.

Do not allow the product to enter any drain during or after application.

Do not apply directly to impervious horizontal surfaces such as sidewalks, driveways, and patios except as a spot or crack-and-crevice treatment.

Treat surfaces to ensure thorough coverage but avoid runoff.

To treat insects harbored in voids and cracks-and-crevices, applications must be made in such a manner to limit dripping and avoid runoff onto untreated structural surfaces and plants.

Do not apply or irrigate to the point of runoff. Do not make applications during rain. Avoid making applications when rainfall is expected before the product has sufficient time to dry (minimum 4 hours).

Rainfall within 24 hours after application may cause unintended runoff of pesticide application.

## STORED GRAIN

Use to control adults and accessible larval stages of listed insect pests including, lesser grain borers, larger grain borers, Angoumois grain moths, Indian meal moths, tobacco moths, Dermestids, and bean weevils; also, to control populations of granary weevils, rice weevils, maize weevils, sawtoothed grain beetles, red flour beetles, and confused flour beetles. Do not apply this product more than once every 30 days.

### Tank-mixing

**Gravista**® may be tank-mixed with other pesticide products such as Pyronyl™ Crop Spray. It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and Directions for Use on all product labels involved in tank-mixing. Users must follow the most restrictive Directions for Use and Precautionary Statements of each product in the tank mixture.

### Directions for Treatment

Dilute **Gravista**® with water or approved dust-controlling oils, mineral or soybean oils and apply to the moving grain stream as a coarse spray.

Final spray volume with water solutions is 3–5 gallons of liquid per 1,000 bushels of grain. For rough rice, use 10 gallons of water per 1,000 bushels. Maintain agitation before and during application.

Use a mixed dilution within 48 hours of preparing. Reagitation of spray dilution may be required if left overnight.

For final spray volumes with oil solutions, follow the oil manufacturer's rate recommendations.

For uniform application, treat the grain as it moves into storage.

Apply **Gravista**® in solution at the rate of 3–5 gallons of dilution per 1,000 bushels to achieve a 0.5 ppm concentration of deltamethrin and 1.2 ppm of S-methoprene on the commodity.

Apply at rates as shown in the following tables.

### AMOUNT OF GRAVISTA®/1,000 BUSHELS OF GRAIN

Commodity	TREATMENT AT 0.5 ppm Deltamethrin/14 ppm PBO/ 1.2 ppm S-Methoprene		Gallons of Diluent	Liters of Diluent
	fl oz	mL		
Barley	31	916	3 - 5	11.3 - 19
Corn	35.6	1,053	3 - 5	11.3 - 19
Oats	20.38	603	3 - 5	11.3 - 19
Popcorn	35.6	1,053	3 - 5	11.3 - 19
Rice	28.5	843	10	11.3 - 19
Rye	35.6	1,053	3 - 5	11.3 - 19
Sorghum (milo)	31.7	937	3 - 5	11.3 - 19
Wheat	38	1,124	3 - 5	11.3 - 19

## AMOUNT OF Gravista®/TON OF GRAIN

Commodity	TREATMENT AT 0.5 ppm Deltamethrin/14 ppm PBO/1.2 ppm S-Methoprene		Ounces of Diluent	Milliliters of Diluent
	fl oz	mL	fl oz	mL
Barley	1.3	38	13 – 21	384 – 630
Corn	1.3	38	13 – 21	384 – 630
Oats	1.3	38	13 – 21	384 – 630
Popcorn	1.3	38	13 – 21	384 – 630
Rice	1.3	38	13 – 21	384 – 630
Rye	1.3	38	13 – 21	384 – 630
Sorghum (milo)	1.3	38	13 – 21	384 – 630
Wheat	1.3	38	13 – 21	384 – 630

### SEED TREATMENT

Apply **Gravista®** to protect stored grain seed against injury from listed insect pests. Clean and treat seed storage areas prior to storage. If seed is infested, fumigate\*\* to control existing populations and apply **Gravista®** for residual protection. **Gravista®** may also be used in conjunction with registered fungicides or other insecticide seed protectants, to protect barley, corn, oats, popcorn, rice, rye, sorghum and wheat seed during the seed treating process, and prior to bagging or placing in bulk seed storage. When treating seed with other pesticide products, properly tag the treated seed, indicating "Do not use seed for food, feed, or oil purposes." Seed treated with **Gravista®** alone is considered grain.

\*\*Fumigate with EPA-registered fumigants according to manufacturer's label recommendations.

### GRAIN BINS AND WAREHOUSES

Dilute 1–6 fl oz of **Gravista®** with water to make 1 gallon of solution. Before storing or handling grain, apply finished spray to equipment, wall and floor surfaces of grain bins and warehouses at the rate of 1 gallon per 1,000 sq ft but do not allow runoff to occur. Apply after equipment bins and buildings have been thoroughly cleaned. If infestations are severe or there are areas that are difficult to treat with a conventional spray, fumigation is recommended with an EPA-registered fumigant.

Use **Gravista®** as a perimeter\* treatment to prevent infestation of these structures. Apply to areas where listed pests are seen or may find entrance. Treat a band of soil or grass up to 6–7 feet wide around and adjacent to structures and treat the foundation to a height of 2 feet. Mix 1–6 fl oz of **Gravista®** with water to make 1 gallon of solution. Apply as a coarse spray at a rate of 1 gallon per 1,000 sq ft. Wet the barrier area thoroughly and uniformly but do not allow runoff to occur. All outdoor\* applications to impervious surfaces are limited to

spot and crack-and-crevice applications only.

\* Not for use in California

**STORAGE AND DISPOSAL:** Do not contaminate water, food, or feed by storage or disposal. **PESTICIDE STORAGE** – Mix as needed. Store in original container in a secured, dry storage area. Avoid extremes in temperatures. Prevent cross-contamination with other pesticides and fertilizers. If the container is leaking and/or material is spilled, on floor or paved surfaces, absorb on sawdust or other commercially available absorbing material, sweep up and remove to chemical waste area for your disposal. Concentrate is stable at normal storage temperatures. **PESTICIDE DISPOSAL** - Wastes resulting from use of this product may be disposed of on site or at an approved waste disposal facility. **CONTAINER HANDLING** – Nonrefillable container. Do not reuse or refill this container. Triple rinse (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then, offer for recycling if available or reconditioning if appropriate or puncture and dispose of in a sanitary landfill or by other procedures approved by state and local authorities.

### IMPORTANT: READ BEFORE USE

Read the entire Directions for Use, Conditions, Disclaimer of Warranties and Limitations of Liability before using this product. If terms are not acceptable, return the unopened product container at once. By using this product, user or buyer accepts the following Conditions, Disclaimer of Warranties and Limitations of Liability.

**CONDITIONS:** The Directions for Use of this product are believed to be adequate and must be followed carefully. However, because of manner of use and other factors beyond Central Garden & Pet Company's control, it is impossible for Central Garden & Pet Company to eliminate all risks associated with the use of this product. As a result, crop injury or ineffectiveness is always possible. All such risks shall be assumed by the user or buyer.

**DISCLAIMER OF WARRANTIES:** To the extent consistent with applicable law, CENTRAL GARDEN & PET COMPANY MAKES NO OTHER WARRANTIES, EXPRESS OR IMPLIED, OR MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE OR OTHERWISE, THAT EXTEND BEYOND THE STATEMENTS MADE ON THIS LABEL. No agent of Central Garden & Pet Company is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. To the extent consistent with applicable law, Central Garden & Pet Company disclaims any liability whatsoever for special, incidental or consequential damages resulting from the use or handling of this product.

**LIMITATIONS OF LIABILITY:** TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER IN CONTRACT, WARRANTY, TORT, NEGLIGENCE, STRICT LIABILITY OR OTHERWISE, SHALL NOT EXCEED THE PURCHASE PRICE PAID, OR AT CENTRAL GARDEN & PET COMPANY'S ELECTION, THE REPLACEMENT OF PRODUCT.

Central Garden & Pet Company  
1501 East Woodfield Road 200W  
Schaumburg, Illinois 60173

Gravista and Pyronyl are trademarks of  
Wellmark International.  
©2024 Wellmark International.

January, 2024  
Schaumburg, IL

# Gravista-D<sup>®</sup>

## Insecticide

**TO CONTROL STORED PRODUCT PESTS IN STORED GRAIN SORGHUM (MILO), CORN, POPCORN, BARLEY, OATS, RICE, RYE, WHEAT**

**PROTECTS STORED GRAINS AND SEEDS AGAINST DAMAGE FROM INDIAN MEAL MOTH, SAWTOOTHED GRAIN BEETLE, RED FLOUR BEETLE, RICE WEEVIL, MAIZE WEEVIL, ALMOND MOTH AND OTHER STORED PRODUCT PESTS**

- Kills stored product adult pests and exposed larvae
- Breaks the pest life cycle and prevents larvae from maturing into adults
- Two modes of action plus a synergist
- Residual control
- For use as an empty bin treatment and direct treatment to stored grain and seeds

### ACTIVE INGREDIENTS:

(S)-Methoprene .....	0.64%
Deltamethrin .....	0.26%
Piperonyl butoxide .....	7.47%
<b>OTHER INGREDIENTS:</b> .....	91.63%
	<u>100.00%</u>

## SPECIMEN LABEL

EPA Reg. No. 89459-123

EPA Est. No. 2724-TX-1

### KEEP OUT OF REACH OF CHILDREN CAUTION

See additional Precautionary Statements and First Aid

Si usted no entiende la etiqueta, busque a alguien para que la explique a usted en detalle.  
(If you do not understand the label, find someone to explain it to you in detail.)

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For general information about this product or emergency medical treatment information, call 1-800-248-7763.

### PRECAUTIONARY STATEMENTS – HAZARDS TO HUMANS AND DOMESTIC ANIMALS – CAUTION

Harmful if inhaled or absorbed through the skin. Causes moderate eye irritation. Avoid breathing dust. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet. Remove and wash contaminated clothing before reuse.

**Personal Protective Equipment (PPE):** Handlers and applicators must wear long-sleeved shirt and long pants, socks and shoes, waterproof gloves, protective eyewear and a PF 10 respirator.

### ENVIRONMENTAL HAZARDS

Do not apply directly to water or to areas where surface water is present or to intertidal areas below the mean high-water mark.

### PRODUCT INFORMATION

GRAVISTA<sup>®</sup>-D INSECTICIDE (GRAVISTA<sup>®</sup>-D) is a ready-to-use formulation that can treat grain as it goes into a storage facility, be applied to commodities prior to storage, or used as an empty bin or silo treatment to areas where commodities will be stored. The residual activity of GRAVISTA<sup>®</sup>-D prevents the emergence of stored product insects and kills exposed adults and insect larvae during storage of commodities. (S)-methoprene, the insect growth regulator in GRAVISTA<sup>®</sup>-D protects the stored products from damage-causing insects by interfering with the normal process of insect development.

The adulticide and synergist work by killing adults and larvae alike. The active ingredients in GRAVISTA<sup>®</sup>-D have established USEPA and CODEX food tolerances [MRLs] for the listed commodities.

GRAVISTA<sup>®</sup>-D is a fine granular insecticide that contains an adulticide, synergist and Insect Growth Regulator. Use to treat grains and empty bins for stored product pest control. GRAVISTA<sup>®</sup>-D provides effective knockdown and control of listed adults and larvae and prevents the buildup of stored product pests with residual control. For use on GRAIN SORGHUM (MILO), CORN, POPCORN, BARLEY, OATS, RICE, RYE, AND WHEAT ONLY.

Use to control adults and accessible larval stages of the following insect pests including, lesser grain borers, larger grain borers, Angoumois

### FIRST AID

**If inhaled** • Move person to fresh air. • **If person is not breathing**, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. • Call a poison control center or doctor for treatment advice.

**If on skin or clothing** • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.

**If in eyes** • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eyes. • Call a poison control center or doctor for treatment advice.

grain moths, Indian meal moths, tobacco moths, Dermestids, and bean weevils; also, to control populations of sawtoothed grain beetles, red flour beetles, confused flour beetles and other stored grain pests.

Granary weevils, rice weevils, maize weevils can be challenging pests to control due to their unique life cycles. In some instances, for these pests, suppression only may be achieved with this product. It is important to monitor for weevil infestations to ensure that they do not exceed the economic threshold.

### DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

**APPLICATION INSTRUCTIONS FOR TREATMENT OF COMMODITIES BEING LOADED OR TURNED INTO STORAGE FACILITIES:** For protection of stored grains and seeds against stored product insects - for optimum results, thoroughly clean and treat storage areas prior to storing commodities. GRAVISTA®-D may be used in conjunction with aeration or an EPA registered fumigation product approved for such use as commonly practiced for grain storage management. Follow the most restrictive of the labeling limitations and precautions.

Apply GRAVISTA®-D to commodities as they are being loaded or turned into storage facilities. Apply 11 pounds of GRAVISTA®-D per 1,000 bushels of commodity or 0.4 pounds (6.4 oz) (181 gms) per ton. Apply the product as uniformly as possible to the grain stream to assure even coverage. Depending on the amount of commodity to be treated, a calibrated hand scoop with the appropriate amount of product may be used to apply into hoppers or boots as the commodity is being moved through an auger into a storage compartment. In larger quantity applications, a powered hopper applicator that applies a metered amount of product to the grain stream, boot or auger may be used. Apply in such a manner as to reduce the potential of the product blowing off target in windy conditions.

**APPLICATION INSTRUCTIONS FOR EMPTY BIN, WALL VOID, CRACK AND CREVICE TREATMENT:** Use GRAVISTA®-D to treat bins, feed mills, and other areas where stored product pests may harbor. For applications to hard surfaces within these sites, apply GRAVISTA®-D at a rate of 13 oz (375 grams) per 1,000 ft<sup>2</sup> of surface area. Pay particular attention to cracks, crevices, and voids where food commodities can accumulate and attract pests. Use a bulb duster or other suitable equipment to apply the material.

**SEED TREATMENT FOR BULK STORAGE:** Apply GRAVISTA®-D to protect grain seed against injury from stored grain insects. Clean and treat seed storage areas prior to loading. Any commercial or on farm treatment equipment for treating commodity as seed is acceptable. Treat using Batch or Continuous flow type equipment. If seed is infested, using the most restrictive of the labeling limitations and restrictions, fumigate\* to control existing populations and apply GRAVISTA®-D for residual protection. Using the most restrictive of the labeling limitations and restrictions, GRAVISTA®-D may also be used in conjunction with registered fungicides or other insecticide seed protectants registered with the EPA, to protect barley, corn, oats, popcorn, rice, rye, sorghum and wheat seed during the seed treating process, and prior to bagging or placing in bulk seed storage. Apply 11 pounds of GRAVISTA®-D per

1,000 bushels of commodity or 0.4 pounds (6.4 oz) (181 gms) per ton. When treating seed with additional pesticide products, properly tag the treated seed, indicating "Treated with (name of substance). Do not use seed for food, feed, or oil purposes." Seed treated with GRAVISTA®-D alone is considered grain. (See below for more information)

\*Using the most restrictive labeling limitations and restrictions, fumigate with EPA-registered fumigants according to manufacturer's label instructions and restrictions.

**SEED BAG LABEL REQUIREMENTS:** Seed commercially treated with this product must be labeled in accordance with all applicable requirements of the Federal Seed Act. The user of this product is responsible for ensuring that the seed bag meets all requirements under the Federal Seed Act.

THE FEDERAL SEED ACT REQUIRES THAT BAGS CONTAINING TREATED SEEDS BE LABELED WITH THE FOLLOWING STATEMENTS:

#### Option 1

This product has been treated with GRAVISTA®-D (89459-123) (S-Methoprene, Deltamethrin, Piperonyl Butoxide) seed treatment. This product as treated is considered to be grain.

#### Option 2

If additional pesticides (i.e., fungicides) are added, list the product name(s) and active(s) as well.

This product has been treated with GRAVISTA®-D (89459-123) (S-Methoprene, Deltamethrin, Piperonyl Butoxide) seed treatment and XXX and YYY products (list out ai's).

– DO NOT use for feed, food, or oil purpose.

The U.S. Environmental Protection Agency requires the following statements on bags containing seeds treated with GRAVISTA®-D seed treatment:

- Store treated seed away from food and feedstuffs.
- Do not allow children, pets, or livestock to have access to treated seeds.
- Treated seeds exposed on soil surface may be hazardous to wildlife. Cover or collect treated seeds spilled during loading and planting (such as in row ends).
- Dispose of seed packaging or containers in accordance with local requirements.

### STORAGE and DISPOSAL

Do not contaminate water, food, or feed by storage or disposal. **Pesticide Storage** - Store in a cool, dry place. **Pesticide Disposal** - Wastes resulting from use of this product must be disposed of on site or at an approved waste disposal facility. **Container Handling** - Nonrefillable container. Do not reuse or refill this container. Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill, or by incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

For information or in case of an emergency, call 1-800-248-7763.

# INSELUX<sup>®</sup>

## FOG & MILL SPRAY

**FOGGING AND SPACE SPRAY FOR USE IN FLOUR MILLS, FEED MILLS, WAREHOUSES, BARNES, FOOD HANDLING ESTABLISHMENTS, AND OTHER STORAGE FACILITIES**

- **Combination insect growth regulator, synergist and adulticide**
- **Kills stored product pest adult insects and exposed larvae** (see Directions for Use for complete listing of pests)
- **Breaks insects' life cycles of stored product pests** (see Directions for Use for complete listing of pests)
- **For use with mechanical, thermal or compressed air fogging systems or space spray**

## SPECIMEN LABEL

### ACTIVE INGREDIENT:

Piperonyl Butoxide*	1.5%
Pyrethrins	0.3%
S-Methoprene	0.3%

<b>OTHER INGREDIENTS:</b>	97.9%
<b>TOTAL</b>	100.0%

\* (butylcarbityl)(6 propylpiperonyl) ether and related compounds

EPA Reg. No. 89459-126

EPA Est. No. 2724-TX-1

**KEEP OUT OF REACH OF CHILDREN**

**CAUTION**

See additional Precautions

### FIRST AID

<b>If on skin or clothing</b>	<ul style="list-style-type: none"> <li>• Take off contaminated clothing.</li> <li>• Rinse skin immediately with plenty of water for 15-20 minutes.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
-------------------------------	---

Have the product container or label with you when calling a poison control center, doctor or going for treatment. You may contact 1-800-248-7763 for information including health concerns, medical emergencies or pesticide incidents.

### PRECAUTIONARY STATEMENTS

#### HAZARDS TO HUMANS AND DOMESTIC ANIMALS

**CAUTION.** Avoid contact with skin or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Wear long-sleeved shirt and long pants, shoes and socks, chemical-resistant gloves.



This mark is not an indication of safety. Read and follow all label instructions.

### User Safety Requirements

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry. Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product. Do not reuse them.

### User Safety Recommendations

Users should wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.

Users should remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

Users should remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible wash thoroughly and change into clean clothing.

### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

### Use Restrictions

Read all directions and cautions before using this product. Apply only as specified on this label.

Remove or cover exposed food and drinking water before application.

Remove or cover dishes, utensils, food processing equipment and food preparation surfaces or wash them before use.

Remove pets and birds and cover fish aquariums before spraying.

Not for use in residential areas.

Do not use in nurseries or rooms where infants, ill or aged persons are confined.

Do not apply directly into sewers or drains or to any area like a gutter where drainage to storm sewers, water bodies or aquatic habitat can occur. Do not allow the product to enter any drain during or after application.

Do not enter or allow others to enter until sprays have dried and vapors, mists and aerosols have dispersed, and the treated area has been thoroughly ventilated for a minimum of 30 minutes.

Keep the area closed for at least 30 minutes and thoroughly ventilate before reoccupying.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.

After space spraying, do not remain in treated area. Exit area immediately and remain outside the treated area until aerosols, vapors and/or mists have dispersed.

When used in dairy barns or facilities: Close milk bulk tank lids to prevent contamination from spray and from dead or falling insects. Remove or cover milking utensils before application. Wash teats of animals before milking.

Do not use in metering devices.

The maximum rate for space spray for any use listed on this label = 2 fl oz per 1,000 cu ft.

**INSELUX® FOG & MILL SPRAY** is a concentrated insecticide and IGR that may be used undiluted or diluted, in equipment capable of applying a fine mist. The product may be applied through cold foggers, thermal foggers, backpack misters or other equipment capable of applying a space spray. **INSELUX® FOG & MILL SPRAY** kills adult insects as well as breaks the insects' life cycles with an insect growth regulator.

### APPLICATION DIRECTIONS

When used according to directions for control of almond moths, Angoumois grain moths, boxelder bugs, bran bugs, cadelles, carpet beetles, cheese mites, cheese skippers, chocolate moths, cigarette beetles, clover mites, crickets, confused flour beetles, drugstore beetles, earwigs, flour beetles, foreign grain beetles, fruit flies, grain beetles, grain mites, granary weevils, non-biting gnats, ham mites, house flies, Indian

meal moths, lesser grain borers, mealworms, mosquitoes, Mediterranean flour moths, merchant grain beetles, nuisance ants (excluding carpenter, fire, harvester and pharaoh ants), rice weevils, red flour beetles, rice flour beetles, rusty grain beetles, sawtoothed grain beetles, silverfish, small flying moths, spider beetles, spiders (except brown recluse and black widow spiders), warehouse beetles, and tobacco moths: **INSELUX® FOG & MILL SPRAY** may be applied undiluted or diluted with mineral oil at a 1:1 ratio through mechanical, thermal or compressed air fogging systems, as a fogging mist or space spray in all approved indoor application sites: auditoriums, bakeries, beverage plants, breweries, bottling plants, buildings, cafeterias, canneries, coffee warehouses and processing plants, concession stands, dairies, day care centers, delicatessens, department stores, diners, dumpsters, egg processing plants, factories, feed mills, flour mills, food processing and storage facilities, food processing plants, freight containers, fruit packing sheds, grain bins/elevators and head spaces, grain harvesting and handling equipment, grain mills, granaries, garages, grocery stores, horse barns, hospitals, hotels, indoor-eating establishments, institutions, locker rooms, meat packing plants, milk rooms, motels, nursing homes, office buildings, public buildings, peanut warehouses and processing plants, restaurants, retail stores, rice mills, stores, schools, ships' holds, sports stadiums, stables, storage bins (areas), stored food in multi-walled paper or cloth bags, supermarkets, theaters, trucks, tobacco factories, trains, warehouses, wineries.

**Cold or Thermal-Fogging and Space Spraying:** For use in aerosol generators to kill exposed stages of listed pests. Apply with equipment designed and adjusted to deliver an aerosol mist. **INSELUX® FOG & MILL SPRAY** is effective against pests listed in the table below. Follow the listed use rates for specific pests. **INSELUX® FOG & MILL SPRAY** may be applied undiluted or diluted as long as the appropriate amount of product is used per 1,000 cubic feet. Apply **INSELUX® FOG & MILL SPRAY** with equipment capable of applying an aerosol mist which includes cold foggers, thermal foggers, backpack misters and other equipment capable of applying a space spray. Close windows and doors and shut off ventilating systems. Extinguish all open flames and pilot lights. Apply at the rate of 1 to 2 fluid ounces per 1,000 cubic feet of space. Reference table for proper rate based on the targeted species. Direct spray towards the ceiling and upper corners of the area and behind obstructions. Keep the area closed for at least 30 minutes and thoroughly ventilate before reoccupying.

Use Sites in Food and Non-Food Handling Establishments	For use as a space spray using cold or thermal foggers, backpack misters or other equipment capable of applying a space spray, to kill exposed stages of the species listed below, use the indicated rate per 1,000 cubic feet	Rate of INSELUX® FOG & MILL SPRAY in fluid ounces
auditoriums bakeries beverage plants bottling plants breweries buildings cafeterias canneries coffee warehouses and processing plants concession stands dairies day care centers delicatessens department stores diners dumpsters egg processing plants factories feed mills flour mills food processing and storage facilities food processing plants freight containers fruit packing sheds grain bins/elevators and head spaces grain harvesting and handling equipment garages grain mills granaries grocery stores horse barns hospitals hotels indoor-eating establishments institutions locker rooms meat packing plants milk rooms motels nursing homes office buildings public buildings peanut warehouses and processing plants restaurants retail stores rice mills schools ships' holds sports stadiums stables stores storage bins (areas) stored food in multi-walled paper or cloth bags supermarkets theaters tobacco factories and warehouses trains trucks warehouses wineries	house flies Indian meal moths mosquitoes rice weevils sawtoothed grain beetles	1 fl oz
	almond moths Angoumois grain moths boxelder bugs bran bugs cadelles carpet beetles cheese mites cheese skippers chocolate moths cigarette beetles clover mites crickets confused flour beetles drugstore beetles earwigs flour beetles foreign grain beetles fruit flies grain beetles grain mites granary weevils gnats, non-biting ham mites lesser grain borers mealworms Mediterranean flour moths merchant grain beetles nuisance ants** red flour beetles rice flour beetles rusty grain beetles silverfish small flying moths spider beetles tobacco moths warehouse beetles spiders***	2 fl oz

\*\*Except carpenter, fire, harvester and pharaoh ants

\*\*\*Except black widows and brown recluse

## STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

**Pesticide Storage.** Store product in its original container in a cool, dry place away from heat or open flame in an area that is inaccessible to children and animals. Avoid exposure to extreme temperatures. If container is damaged: Stop any leaks by repositioning the container or by patching or otherwise repairing the leaks. Take care to avoid contact with pesticide and wear protective gear. On cleanup of spilled liquids, wear protective equipment as required to prevent contact with the product or its vapors. Cover the spilled areas with generous amounts of absorbent material such as clay, diatomaceous earth, sand or sawdust. Sweep the contaminated absorbent onto a shovel and put the sweepings into a salvage drum. Dispose of wastes as below.

**Pesticide Disposal.** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. **Container Handling. (5 gallons or less)** Nonrefillable container. Do not reuse or refill this container. Clean container promptly after emptying. Triple rinse (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain container for 10 seconds after the flow begins to drip. Fill the container  $\frac{1}{4}$  full with solvent used to dilute product and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or reconditioning if appropriate or puncture and dispose of in a sanitary landfill or by other procedures approved by state and local authorities.

**(greater than 5 gallons)** Nonrefillable container. Do not reuse or refill this container. Clean container promptly after emptying. Triple rinse (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container  $\frac{1}{4}$  full with solvent used to dilute product. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Then offer for recycling if available or reconditioning if appropriate, or puncture and dispose of in a sanitary landfill or by other procedures approved by state and local authorities.

## WARRANTY AND CONDITIONS OF SALE

To the fullest extent permitted by law, Seller makes no warranty, expressed or implied, concerning the use and handling of this product other than indicated on the label. To the extent permitted by law, Buyer assumes all risks of use and handling of this material when such use and handling are contrary to label instructions.

For information or in case of an emergency, call 1-800-248-7763.

Manufactured for:

**Central Garden & Pet Company**  
1501 East Woodfield Road 200W  
Schaumburg, Illinois 60173

Inselux is a registered trademark of  
Wellmark International.  
©2024 Wellmark International.

January, 2024  
Schaumburg, IL

# PBO-8<sup>®</sup>

## Synergist

**FOR USE IN:** residential dwellings or indoors in commercial or institutional buildings (including food processing and handling areas), turf and ornamentals, crops (including greenhouses and nurseries), on livestock, and livestock housing.

An emulsifiable concentrate containing 8 lbs. of piperonyl butoxide per gallon.

- Condensed specimen label – view full specimen label at [www.bugfreegrains.com](http://www.bugfreegrains.com)

## SPECIMEN LABEL

### ACTIVE INGREDIENTS:

Piperonyl Butoxide\* ..... 91.3%

**OTHER INGREDIENTS:**..... 8.7%

**TOTAL:** ..... 100.0%

\* (butylcarbityl)(6-propylpiperonyl) ether and related compounds.

**KEEP OUT OF REACH OF CHILDREN  
CAUTION**

See Additional Precautionary Statements and Directions for Use

EPA Reg.No. 89459-33  
EPA Est. No. 2724-TX-1

### PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

**CAUTION:** Harmful if swallowed or absorbed through the skin. Avoid contact with skin, eyes or clothing. Wear long-sleeved shirt and long pants, socks and shoes and chemical-resistant gloves (such as barrier laminate, butyl rubber, nitrile rubber, neoprene rubber, polyvinyl chloride or viton (Selection Category C)).

FIRST AID	
<b>If on skin or clothing</b>	<ul style="list-style-type: none"> <li>• Take off contaminated clothing.</li> <li>• Rinse skin immediately with plenty of water for 15-20 minutes.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b>If swallowed</b>	<ul style="list-style-type: none"> <li>• Call a poison control center or doctor immediately for treatment advice.</li> <li>• Have person sip a glass of water if able to swallow.</li> <li>• Do not induce vomiting unless told to do so by a poison control center or doctor.</li> <li>• Do not give anything by mouth to an unconscious or convulsing person.</li> </ul>
<b>If in eyes</b>	<ul style="list-style-type: none"> <li>• Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li> <li>• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul> <p style="text-align: right;"><i>(continued)</i></p>

<b>If inhaled</b>	<ul style="list-style-type: none"> <li>• Move person to fresh air.</li> <li>• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible.</li> <li>• Call a poison control center or doctor for further treatment advice.</li> </ul>
-------------------	---

**HOT LINE NUMBER:** Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-347-8272 for emergency medical treatment information.

### Personal Protective Equipment (PPE):

Mixers, loaders, applicators and other handlers must wear the following: long sleeved shirt and long pants, shoes and socks and chemical-resistant gloves.

In addition to the above PPE, mixers and loaders supporting aerial applications to forests and/or applicators using a high pressure hand wand in an enclosed area must wear at least a NIOSH-approved respirator with a dust/mist filter with MSHA/NIOSH approval number prefix TC-21C or any N, R, P or HE filter.

In addition to the above PPE, applicators using hand held foggers in an enclosed area must wear a half-face, full-face, or hood-style NIOSH-approved respirator with a dust/mist filtering cartridge (MSHA/NIOSH approval number prefix TC-21C), or a canister approved for pesticides (MSHA/NIOSH approval number prefix TC-14G), or a cartridge or canister with any N, R, P or HE filter. See Engineering Controls for additional requirements.

**User Safety Requirements:** Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry. Discard

clothing and other absorbent materials that have been drenched or heavily contaminated with the product's concentrate. Do not reuse them.

**Engineering Controls:** Pilots must use an enclosed cockpit that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(6)]. Human flagging is prohibited. Flagging to support aerial application is limited to use of the Global Positioning System (GPS) or mechanical flaggers.

**User Safety Recommendations:**

Users should wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Users should remove clothing/PPE immediately if pesticide gets inside, then wash thoroughly and put on clean clothing. Users should remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

**ENVIRONMENTAL HAZARDS**

This product is toxic to aquatic organisms, including fish and invertebrates. Drift and run-off may be hazardous to aquatic organisms in water adjacent to treated areas. This product has the potential to contaminate water through run-off. This product has a potential for run-off for several weeks after application. Poorly draining soils and soils with shallow water tables are more prone to produce run-off that contains this product.

Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters or rinsate.

Do not discharge effluent containing this product into lakes, streams, ponds, estuaries, oceans, or other waters unless in accordance with the requirements of a National Pollution Discharge Elimination System (NPDES) permit, and the permitting authority has been notified in writing prior to discharge. Do not discharge effluent containing this product to sewer systems without previously notifying the local sewage treatment plant authority. For guidance, contact your State Water Board or Regional Office of the EPA.

**PHYSICAL AND CHEMICAL HAZARDS**

Do not use or store near heat or open flame.

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

**USE RESTRICTIONS**

This product is an emulsifiable concentrate for use as an additive in insecticides and acaricides to improve control of insects and mites. This product aids in reducing oxidative resistance that can develop in insects and mites. This product has no insecticidal effect of its own.

- Apply this product only as specified on this label.
- Do not contaminate food or feedstuffs.
- Except when applying to livestock, dairy and poultry animals, do not apply this product in a way that will contact workers or other persons, either directly or through drift.
- Except when applying to livestock, dairy and poultry animals, only protected handlers may be in the treatment area during application. Do not enter or allow others to enter the treated area until vapors, mists, and aerosols have dispersed, and the treated area has been thoroughly ventilated.
- Not for use in outdoor residential misting systems (indoor or outdoor). Do not use in nurseries or rooms where infants, ill or aged persons are confined. Do not place metering device directly over or within 8 feet of exposed food, dishes, utensils, food processing equipment and food handling or preparation areas. If the device works on an automatic timer, set the timing for application when people, pets, and/or food are unlikely to be present. If the device works when a person operates a remote activation

*(continued)*

device, then application of this pesticide when people, pets, and food are present is prohibited. Direct nozzle to spray towards the target area and away from areas where people are typically present. The pesticide container must be locked. Do not install within 3 feet of air vents, air conditioner units, or windows. Carefully follow directions for the dispenser unit when installing the dispenser and replacing cans or conducting maintenance.

- Do not use in aircraft cabins.
- Do not wet articles to point of run-off or drip. Do not use treated articles until spray has dried.
- Do not wet plants to point of run-off or drip.
- Do not apply this product directly into sewers or drains, or to any area like a gutter where drainage to storm sewers, storm drains, water bodies, or aquatic habitat can occur. Do not allow this product to enter any drain during or after application.
- Do not make applications when heavy rain is expected or during rain.
- Not for use in outdoor residential misting systems.
- Do not remain in treated area. Exit area immediately and remain outside the treated area until aerosols, vapors and/or mists have dispersed.
- Remove or cover exposed food and drinking water before application. Remove or cover dishes, utensils, food processing equipment and food preparation surfaces, or wash them before use.
- When used in dairy barns or facilities: Close milk bulk tank lids to prevent contamination from spray and from dead or falling insects. Remove or cover milking utensils before application. Wash teats of animals before milking.
- For agricultural premises, commercial animal housing and equipment, direct application to non-domestic animals/livestock, greenhouse grown ornamental, flowering and foliage plants and outdoor ornamentals and lawns: Do not apply more than 1 time per day.
- For food crops grown outdoors or in greenhouses in agricultural settings: Do not apply more than 10 times per season. Do not reapply within 3 days except under extreme pest pressure. In case of extreme pest pressure, do not reapply within 24 hours. Do not apply to cotton within 14 days of seed harvest.
- For indoor food handling/processing facilities: Do not make space spray applications when facility is in operation. Prior to space spray applications, cover or remove food. Prior to space spray applications, cover food processing surfaces or clean after treatment and before use. Do not apply more than 1 time per day.
- For post-harvest application to vegetables, fruits and nuts: Do not apply more than 1 time per day. Do not reapply within 7 days. Do not apply more than 10 times to sweet potatoes.
- For post-harvest application to stored grain and seed: Do not reapply within 30 days.
- For food stored in bags: Direct application to food contact surfaces is prohibited.
- For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.
- Remove pets, birds, and cover fish aquariums before spraying.

**AGRICULTURAL USE REQUIREMENTS**

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR, Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

*(continued)*

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours. PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is: coveralls, chemical-resistant gloves, such as barrier laminate, or butyl rubber, or nitrile rubber, or neoprene rubber, or PVC, or viton, and shoes plus socks.

### NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR, Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

**For surface spray use:** Except when applying to livestock, dairy and poultry animals, do not enter or allow others to enter treated area until sprays have dried.

**For space spray use:** Do not enter or allow others to enter until vapors, mists, or aerosols have dispersed and the treated area has been thoroughly ventilated.

**TANK MIXES:** Tank mixing or use of this product with any other product which is not specifically and expressly authorized by the label shall be the exclusive risk of user, applicator, and/or application advisor. Read and follow the most restrictive label instructions and precautions of all products in a tank mix.

### APPLICATION INSTRUCTIONS

This product may be applied using ground equipment sprayers, hand-held sprayers, power sprayers, foggers, ULV applicators, or by chemigation.

Not for use in outdoor residential misting systems.

### NON-CROP USES

This product may be mixed with insecticides registered for use in the following areas: Residential dwellings or indoors in commercial or institutional buildings including houses (indoor or outdoor), multiple unit housing, hotels, schools, restaurants, food processing facilities, food storage facilities, bakeries, healthcare facilities, warehouses, office buildings, factories, industrial facilities. Do not use in aircraft cabins.

For correct rate of insecticide, refer to the label of the tank-mix insecticide.

When used as a space spray, do not exceed 0.4 fl oz of this product per 1,000 cu ft of treated area. Do not apply more than 1 time per day. Thoroughly ventilate the treated area before re-entry or reoccupying.

If used as a crack and crevice or broadcast surface spray, do not exceed 8 fl oz of this product per 1,000 sq ft of treated area. Do not apply more than 1 time per day.

For metered-release device space sprays, do not exceed 0.8 fl oz of this product per 1,000 cu ft of treated area. Do not apply more than 1 time per day.

### FOOD HANDLING ESTABLISHMENTS

(Places other than private residences in which food is held, processed, prepared or served)

**NON-FOOD AREAS:** This product may be used to treat non-food areas such as garbage rooms, lavatories, entries and vestibules, offices, locker rooms, machine rooms, boiler rooms, garages, mop closets and storage (after canning or bottling).

**FOOD AREAS:** This product can be used to treat food areas of food handling establishments, restaurants or other areas where foods are commercially prepared or served.

### POST-HARVEST APPLICATION TO FOOD COMMODITIES

Do not apply this product except in mixture with an insecticide registered for use in post-harvest application to food commodities. Carefully follow label directions for the insecticide being used. This product may be added to appropriately labeled insecticides on: potatoes, tomatoes, peas, sweet potatoes, apples, blackberries, blueberries, boysenberries, cherries, crabapples, currants, dewberries, figs, gooseberries, grapes, guava, loganberries, mangoes, muskmelons, oranges, peaches, pears, pineapples, plums, raspberries, almonds, walnuts, copra and peanuts.

When used as a space spray to fruits, vegetables and copra, do not exceed 0.16 fl oz of this product per 1,000 cu ft of space treated. When used as a surface application to the commodities listed, do not exceed 1.6 fl oz of product per 1,000 sq ft of treated area. When used as a surface application to fruit or tomatoes in baskets or hampers, do not exceed 0.013 fl oz of this product per 1,000 lb of fruit. When used as a dip or spray, do not exceed 0.64 fl oz of this product per 10 gallons of water.

### STORED GRAIN AND SEED

Maximum rates are listed below. Do not reapply within 30 days.

barley	0.53 fl oz per 100 lb commodity
beans	0.43 fl oz per 100 lb commodity
lima beans	0.46 fl oz per 100 lb commodity
birdseed	0.51 fl oz per 100 lb commodity
buckwheat	0.53 fl oz per 100 lb commodity
cocoa beans	0.59 fl oz per 100 lb commodity
corn	0.46 fl oz per 100 lb commodity
cottonseed	0.91 fl oz per 100 lb commodity
flax	0.46 fl oz per 100 lb commodity
grain sorghum	0.46 fl oz per 100 lb commodity
oats	0.80 fl oz per 100 lb commodity
field peas	0.43 fl oz per 100 lb commodity
rice	0.58 fl oz per 100 lb commodity
rye	0.46 fl oz per 100 lb commodity
wheat (club, common, durum) (emmer, spelt)	0.43 fl oz per 100 lb commodity 0.64 fl oz per 100 lb commodity

### FOOD STORED IN BAGS

Storage areas containing food in impermeable bags may be treated with a space spray. Do not exceed 0.4 fl oz of this product per 1,000 cu ft of space treated. Direct application to food contact surfaces is prohibited.

### OUTDOOR AGRICULTURAL PREMISES

When used as a broadcast surface spray application, do not exceed 0.32 fl oz of this product per 1,000 sq ft treated (16 fl oz per acre). For crack and crevice or spot surface sprays (including applications for fire ants or turf diagnostic aid), do not exceed 8 fl oz of this product per 1,000 sq ft of space treated.

Do not apply more than 1 time per day.

## STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal. Open dumping is prohibited.

**PESTICIDE STORAGE:** If container is damaged: Stop leaks by repositioning the container or by patching or otherwise repairing the leaks. Take care to avoid contact with the pesticide and wear protective gear. On clean up of spilled liquids, wear protective equipment as required to prevent contact with the product or its vapors. Cover the spilled areas with generous amounts of absorbent material, such as clay, diatomaceous earth, sand or sawdust. Sweep the contaminated absorbent onto a shovel and put the sweepings into a salvage drum. Dispose of wastes as below. Place any leaking container into a similar drum or glass container. Do not store, use, pour or spill near heat or open flames.

**PESTICIDE DISPOSAL:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

**CONTAINER DISPOSAL:** Nonrefillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then, offer for recycling if available or reconditioning if appropriate or puncture and dispose of in a sanitary landfill or by other procedures approved by state and local authorities.

## CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

**NOTICE:** Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this Product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather, presence of other materials or other influencing factors in the use of the product, which are beyond the control of Central Garden & Pet Company or Seller. To the extent consistent with applicable law, all such risks shall be assumed by Buyer and User, and Buyer and User agree to hold Central Garden & Pet Company and Seller harmless for any claims relating to such factors.

Central Garden & Pet Company warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. This warranty does not extend to the use of this product contrary to label instructions, or under abnormal conditions or under conditions not reasonably foreseeable to or beyond the control of Seller or Central Garden & Pet Company, and Buyer and User assume the risk of any such use. To the extent consistent with applicable law, Central Garden & Pet Company MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.

To the extent consistent with applicable law, Central Garden & Pet Company or Seller shall not be liable for any incidental, consequential or special damages resulting from the use or handling of this product. To the extent consistent with applicable law, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF CENTRAL GARDEN & PET COMPANY AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF CENTRAL GARDEN & PET COMPANY OR SELLER, THE REPLACEMENT OF THE PRODUCT.

Central Garden & Pet Company and Seller offer this product, and Buyer and User accept it, subject to the foregoing Conditions of Sale and Limitation of Warranty and Liability, which may not be modified except by written agreement signed by a duly authorized representative of Central Garden & Pet Company.

Pyronyl™

# Crop Spray

**KILLS:** Ants, Aphids, Bed Bugs, Black Widow Spiders, Cockroaches, Crickets, Face Flies, Fleas, Horse Flies, House Flies, Lice, Mosquitoes, Stable Flies, Stored Product Insects (including Almond Moths, Confused Flour Beetles, Granary Weevils, Indian Meal Moths, Lesser Grain Borers, Rice Weevils, Saw-toothed Grain Beetles and others listed on this label) Ticks, and Thrips

**FOR USE AS:** Stored Product Protection, Livestock and Poultry Spray

Refer to the label for additional use sites.

This Specimen Label is for Stored Grain Protection. View full label at [www.bugfreegrains.com](http://www.bugfreegrains.com)

**ACTIVE INGREDIENTS:**

Pyrethrins . . . . . 6%

Piperonyl Butoxide\* . . . . . 60%

**OTHER INGREDIENTS\*\*:** . . . . . 34%

TOTAL . . . . . 100%

\*(butylcarbityl)(6-propylpiperonyl) ether and related compounds.

\*\*Contains Petroleum Distillates

EPA Reg. No. 89459-26

EPA Est. No. 2724-TX-1

## SPECIMEN LABEL

**KEEP OUT OF REACH OF CHILDREN**

**CAUTION**

See additional Precautionary Statements,  
First Aid Statements and Directions for Use

### PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

**CAUTION:** Harmful if swallowed. Causes moderate eye irritation. Avoid contact with skin, eyes or clothing. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals.

#### FIRST AID

<b>If swallowed</b>	<ul style="list-style-type: none"> <li>• Immediately call a Poison Control Center or physician.</li> <li>• Do not induce vomiting unless told to do so by a Poison Control Center or physician.</li> <li>• Do not give <b>any</b> liquid to the person.</li> <li>• Do not give anything by mouth to an unconscious person.</li> </ul>
<b>If on skin or clothing</b>	<ul style="list-style-type: none"> <li>• Take off contaminated clothing.</li> <li>• Rinse skin immediately with plenty of water for 15-20 minutes.</li> <li>• Call a Poison Control Center or physician for treatment advice.</li> </ul>
<b>If in eyes</b>	<ul style="list-style-type: none"> <li>• Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li> <li>• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>• Call a Poison Control Center or physician for treatment advice.</li> </ul>

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may contact 1-800-248-7763 for information including health concerns, medical emergencies or pesticide incidents.

**NOTE TO PHYSICIAN:** Contains petroleum distillate – vomiting may cause aspiration pneumonia.

**Personal Protective Equipment (PPE):** Some materials that are chemical-resistant to this product are made of barrier laminate, nitrile rubber, neoprene rubber, viton.

Mixers, loaders, applicators, and other handlers must wear the following:

- long-sleeve shirt,
- long pants,
- shoes and socks and
- chemical-resistant gloves.

In addition to the above PPE, applicators using a high pressure handwand in an enclosed area must wear at least a NIOSH-approved respirator with:

- a filter with NIOSH approval number prefix TC-21C or
- any R, P, or HE filter.

In addition to the above PPE, applicators using hand held foggers in an enclosed area must wear a half-face, full-face, or hood-style NIOSH-approved respirator with:

- a filtering cartridge (NIOSH approval number prefix TC-21C), or
- a canister (NIOSH approval number prefix TC-14G), or
- a cartridge or canister with any R, P, or HE filter.

See Engineering Controls for additional requirements.

**User Safety Requirements:** Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with the product's concentrate. Do not reuse them.

**User Safety Recommendations:** Users should wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Users should remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

Users should remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

## ENVIRONMENTAL HAZARDS

**For terrestrial applications:** This product is toxic to aquatic organisms, including fish and invertebrates. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. This product may contaminate water through runoff. This product has potential for runoff for several weeks after application. Poorly draining soils and soils with shallow water tables are more prone to produce runoff that contains this product.

This product is highly toxic to bees exposed to direct treatment on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds while bees are foraging the treatment area.

Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters or rinsate.

## DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. For any requirements specific to your state or tribe, consult the agency in your state responsible for pesticide regulations.

### General Precautions & Use Restrictions:

-When applied to outdoor residential surfaces:

All outdoor applications must be limited to spot or crack-crevice treatments only, except for the following permitted uses: 1) Treatment to soil or vegetation around structures; 2) Applications to lawns, turf and other vegetation; 3) Applications to building foundations, up to a maximum height of 3 feet.

Other than applications to building foundations, all outdoor applications to impervious surfaces such as sidewalks, driveways, patios, porches and structural surfaces (such as windows, doors and eaves) are limited to spot and crack-and-crevice applications only.

-For surface treatment of stored grain and seed: Do not reapply within 30 days.

## AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and the handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is: coveralls, chemical-resistant gloves made of any waterproof material and shoes plus socks.

## NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

### For surface spray use:

- Except when applying to livestock and as a mosquito adulticide, do not enter or allow others to enter treated area until sprays have dried.

### For space spray use:

- Except when applying as a mosquito adulticide, do not enter or allow others to enter until vapors, mists and aerosols have dispersed and the treated area has been thoroughly ventilated.

## USE AS A SURFACE SPRAY

**IN HOMES, RESTAURANTS, FOOD PROCESSING PLANTS, INDUSTRIAL INSTALLATIONS AND WAREHOUSES:** To kill accessible, exposed stages of crawling insects including, but not limited to, ants, cockroaches, cadelles, cigarette beetles, confused flour beetles, dark mealworms, dried fruit beetles, drugstore beetles, grain mites, red flour beetles, rice weevils, saw-toothed grain beetles, spider beetles, yellow mealworms, dilute 1 part Pyronyl™ Crop Spray with 59 parts water and apply at the

rate of 1 gallon to 750 square feet, paying special attention to force the spray into all cracks and crevices.

**IN COMBINATION WITH RESIDUAL INSECTICIDES:** To provide flushing and quick knockdown of insects, this product may be tank-mixed with other insecticides at the rate of ¼ to ½ fluid ounce (equivalent to ½ to 1 tablespoon or 7.4 ml. to 14.8 ml.) per gallon of finished spray.

**IN USDA INSPECTED FACILITIES:** To kill accessible, exposed stages of crawling insects including, but not limited to ants, cockroaches, cadelles, cigarette beetles, confused flour beetles, dark mealworms, dried fruit beetles, drugstore beetles, grain mites, red flour beetles, rice weevils, saw-toothed grain beetles, spider beetles, yellow mealworms, dilute 1 part of Pyronyl™ Crop Spray with 19 parts of water and apply at the rate of 1 gallon to 750 square feet, paying special attention to force the spray into all cracks and crevices.

## USE AS A SPACE SPRAY

To kill crawling and flying insects in sites that include homes, restaurants, food processing plants, industrial installations and warehouses, dilute Pyronyl™ Crop Spray with water and apply as a space spray. For best results, close doors and windows before spraying and keep them closed for 30 minutes after treatment. Where oil residues are not undesirable, Pyronyl™ Crop Spray can be diluted in deodorized base oil instead of water and applied with mechanical, thermal or ULV applicators.

**CRAWLING AND FLYING INSECTS:** To kill accessible, exposed stages of CRAWLING INSECTS including ants, cockroaches, cadelles, cigarette beetles, confused flour beetles, dark mealworms, dried fruit beetles, drugstore beetles, grain mites, red flour beetles, rice weevils, saw-toothed grain beetles, spider beetles, yellow mealworms and FLYING INSECTS including Angoumois grain moths, Indian meal moths, mosquitoes, Mediterranean flour moths, small flying moths, tobacco moths, add 10.67 fl oz (1 part to 11 parts water or oil) of Pyronyl™ Crop Spray per gallon of oil or water and apply at the rate of 1 fluid ounce per 1,000 cubic feet of space. Direct the spray towards the ceiling and upper corners of the area and behind obstructions. Keep the area closed for at least 30 minutes after treatment.

**FLYING INSECTS:** To kill flying insects including Angoumois grain moths, cheese skippers, fruit flies, fungus gnats, gnats, house flies, Indian meal moths, mosquitoes, Mediterranean flour moths, small flying moths, tobacco moths, dilute 1 part of Pyronyl™ Crop Spray with 47 parts of water or oil (2.67 ounces per gallon) and apply at the rate of 1 fluid ounce per 1,000 cubic feet of space. Direct the spray towards the ceiling and upper corners of the area and behind obstructions. Keep the area closed for at least 30 minutes after treatment.

**FOR USE ON SWEET POTATOES IN STORAGE IN COMMERCIAL STORAGE/ WAREHOUSES PREMISES:** To kill fruit flies and vinegar flies, dilute this concentrate at 1 part to 19 parts water (6.4 fluid ounces per gallon (51 ml/L)). Apply as a space fog with a mechanical fogger capable of producing particles of aerosol size at the rate of 1 gallon diluted spray per 100,000 cubic feet (1.34 ml/m<sup>2</sup>) of space. Apply only when flying insects are present. Several applications may be necessary during periods of heavy infestation. Do not reapply within 7 days except under extreme pest pressure. In case of extreme pest pressure, do not reapply within 24 hours. Do not apply more than 10 times to sweet potatoes.

## USE IN STORED PRODUCT PROTECTION

**AS A GRAIN AND SEED PROTECTANT:** Pyronyl™ Crop Spray may be applied to the following grains and seeds: barley, beans, birdseed, buckwheat, cocoa beans, corn, cottonseed, flax, oats, peas (field) rice, rye, sorghum and wheat to protect them from grain storage insects for full a season or approximately 8 months. Pyronyl™ Crop Spray may be used in combination with a registered fumigant for use of heavily infested stored products.

**TO KILL STORED PRODUCT INSECTS INCLUDING** almond moths, Angoumois grain moths, cadelles, cigarette beetles, confused flour beetles, drugstore beetles, flat grain beetles, granary weevils, Indian meal moths, lesser grain borers, maize weevils, Mediterranean flour moths, merchant grain beetles, red flour beetles, rice weevils, rusty grain beetles, saw-toothed grain beetles and squarenecked grain beetles, dilute at the rate of 1 part Pyronyl™ Crop Spray with 29 parts water (1 pint with 3 gallons 5 pints water). Thoroughly mix the emulsion and apply at the rate of 4 to 5 gallons per 1,000 bushels of grain or seed as it is carried along a belt or as it enters the auger or elevator.

**SURFACE TREATMENT OF STORED GRAIN AND SEED:** To kill Indian meal moths, Angoumois grain moths and Mediterranean flour moths, inspect monthly after the grain is placed in storage. If the top 2 or 3 inches are infested, dilute 1 part of

Pyronyl™ Crop Spray with 19 parts of water and apply at the rate of 1 to 2 gallons per 1,000 square feet of grain. Rake the mixture into the grain to a depth of 4 inches.

**ON ALMONDS, PEANUTS AND WALNUTS IN BULK OR IN BAGS:** To kill stored product insects including almond moths, Angoumois grain moths, ants, cadelles, cigarette beetles, confused flour beetles, drugstore beetles, flat gain beetles, granary weevils, Indian meal moths, lesser grain borers, maize weevils, Mediterranean flour moths, merchant grain beetles, red flour beetles, rice weevils, rusty grain beetles, saw-toothed grain beetles and square-necked grain beetles, dilute 1.33 fluid ounces of Pyronyl™ Crop Spray per gallon of water and apply a coarse wet spray over the top of stored nuts or the outer surface of stacked bagged nuts at the rate of 4 gallons per 1,000 square feet. Apply at weekly intervals for about 6 weeks and then at 15 day intervals. Apply the first two applications at the rate of 2 gallons per 1,000 square feet.

**STORAGE SITES:** To treat grain and seed storage sites, warehouse bins, trucks, cargo ships and planes prior to filling with grain or seed, thoroughly clean the site by sweeping out the waste, cobwebs, and other debris on the walls and rafters as well as on floor and about the door frames, paying special attention to the material lodged in the cracks and crevices. Remove these accumulations and burn to kill eggs and insects that might be present.

In mills and elevators, pay particular attention to the bin hoppers to remove all grain infested accumulations. Clean and insure that conveying equipment are free of trash deposits that could maintain an infestation. For farms, clean up and around the used feed and grain bags, grain residues from wagons, harvesting equipment and feed troughs. Do not place newly harvested grain in the same bin with carry-over grain and fumigate all carry-over grain stocks not treated with grain protectant. Perform cleaning operations within two or three weeks before harvest.

To treat the storage site prior to using it for storage, dilute 1 part of Pyronyl™ Crop Spray with 59 parts of water ((1 pint with 7 gallons 3 pints of water) and apply to walls, floors, ceilings and partition boards at the rate of one gallon per 750 sq. feet. It is important to thoroughly treat all crack and crevices.

## STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

**PESTICIDE STORAGE AND SPILL PROCEDURES:** Store upright at room temperature. Avoid exposure to extreme temperatures. In case of spill or leakage soak up with absorbent material such as sand, sawdust, earth, fuller's earth, etc. Dispose of with chemical waste.

**PESTICIDE DISPOSAL:** Pesticide, spray mixture, or rinse water that cannot be used according to label instructions must be disposed of at or by an approved waste disposal facility.

**CONTAINER DISPOSAL:** Non-refillable container. Do not reuse or refill this container. Clean container promptly after emptying. **{Containers 5 gallons or less}** Triple rinse as follows. Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water (or solvent used to dilute product) and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then, offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other approved state and local procedures. **{For containers greater than 5 gallons}** Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container ¼ full with water (or solvent used to dilute product). Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution for 30 seconds. Stand the container on its end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other approved state and local procedures.

[www.bugfreegrains.com](http://www.bugfreegrains.com)

**Central Garden & Pet Company**, 1501 East Woodfield Road, 200W, Schaumburg, Illinois 60173

NOTE: This specimen label is for informational purposes only. All uses may not be approved in all states. See product labeling for use directions. Pyronyl is a trademark of Wellmark International. © 2015, 2017 Wellmark International.







**BUG FREE**  
**GRAINS**  
*Stored Product Solutions*



[www.bugfreegrains.com](http://www.bugfreegrains.com)



Always read and follow label directions. PBO-8, Gravista, Inselux, Diacon, Centynal and Pyronyl are trademarks of Wellmark International. The red to yellow gradation is a registered trademark of Central Garden & Pet Company.  
©2024 Wellmark International. AG 24-002